## Georgia State University Digital Archive @ GSU

**Public Health Theses** 

Institute of Public Health

7-23-2007

### Multiple Sclerosis Disease Distribution and Potential Impact of Environmental Air Pollutants in Georgia

Anthony Charles Gregory

Follow this and additional works at: http://digitalarchive.gsu.edu/iph theses

#### Recommended Citation

Gregory, Anthony Charles, "Multiple Sclerosis Disease Distribution and Potential Impact of Environmental Air Pollutants in Georgia" (2007). *Public Health Theses.* Paper 8.

This Thesis is brought to you for free and open access by the Institute of Public Health at Digital Archive @ GSU. It has been accepted for inclusion in Public Health Theses by an authorized administrator of Digital Archive @ GSU. For more information, please contact digital archive@gsu.edu.

# MULTIPLE SCLEROSIS DISEASE DISTRIBUTION AND POTENTIAL IMPACT OF ${\rm ENVIRONMENTAL~AIR~POLLUTANTS~IN~GEORGIA}$

By

## ANTHONY CHARLES GREGORY II B.S. GEORGIA STATE UNIVERSITY

A Thesis Submitted to the Graduate Faculty of the Georgia State University in Partial
Fulfillment of the Requirements for the Degree
MASTER IN PUBLIC HEALTH
ATLANTA, GEORGIA

#### APPROVAL PAGE

## MULTIPLE SCLEROSIS DISEASE DISTRIBUTION AND POTENTIAL IMPACT $$\operatorname{\textsc{OF}}$$

#### ENVIRONMENTAL EXPOSURES IN GEORGIA

By

Anthony Charles Gregory II

Approved:		
Dr. Derek G. Shendell		
Committee Chair		
Dr. Solomon I. Okosun		
Committee Member		
Dr. Karen. E. Gieseker		
Committee Member		

#### ANTHONY C. GREGORY II

Multiple Sclerosis Disease Distribution and Potential Impact of Environmental Air Pollutants in Georgia.

Under the direction of Derek G. Shendell, D.Env, MPH

This study's purpose is to examine the potential relationships between MS (Multiple Sclerosis) and environmental outdoor air pollutants in GA (Georgia). We used secondary, cross-sectional data received from the member's list of the National Multiples Sclerosis Society's GA chapter (NMSS-GA), the U.S. census, and scorecard.org, a database based on the U.S. Environmental Protection Agency (EPA) Toxic Release Inventory (to 2002) and National Emissions Inventory (1999) data. The study population was 9,072,576 including 6,247 self-reported MS cases from NMSS-GA.

Using stepwise multivariate linear regression of census and EPA data, the best predictive model in GA for female or total prevalence rates used per capita income and coarse particles (PM-10); male prevalence rates included only per capita income.

The relationship between MS and PM-10 could be facilitated by influences of PM-10 on the systemic immune response and inflammation. More exposure and basic and clinical research are needed to understand environmental influences on MS.

INDEX WORDS: Multiple Sclerosis, Environment, Georgia, Epidemiology, Air Pollutants, Toxicants, Particulate Matter

#### **ACKNOWLEDGEMENTS**

I thank my committee members, Dr. Shendell (chair), Dr. Okosun and Dr. Gieseker, for their support throughout this process. Also, I thank Ms. Hilary LaMotte and the National Multiple Sclerosis Society's support of this project. Most of all, I thank my wife, Denise, for her unfailing support and help throughout this process. Without these people supporting this project, this work could not have been done.

V

**AUTHOR'S STATEMENT** 

In presenting this thesis as a partial fulfillment of the requirements for an

advanced degree from Georgia State University, I agree that the Library of the

University shall make it available for inspection and circulation in accordance with its

regulations governing materials of this type. I agree that permissions to quote from,

to copy from, or to publish this thesis may be granted by the author or, in his/her

absence, by the Associate Dean College of Health and Human Sciences. Such

quoting, copying, or publishing must be solely for scholarly purposes and will not

involve potential financial gain. It is understood that any copying from or publication

of this thesis which involves potential financial gain will not be allowed without

written permission of the author.

Anthony C. Gregory II

\_\_\_\_

Signature of the Author.

#### NOTICE TO BORROWERS PAGE

#### **Notice to Borrowers Page**

All theses deposited in the Georgia State University Library must be used in accordance with the stipulations prescribed by the author in the preceding statement.

The author of this thesis is:

Student's Name: Anthony C. Gregory II

Street Address: 6871 Peachtree Dunwoody Rd. NE, Apt 210

City, State, Zip Code: Atlanta, GA, 30328

Professor's Name: Derek Shendell, D. Env, MPH

Department: Institute of Public Health

College: College of Health and Human Sciences

Georgia State University

P.O. Box 4018

Atlanta, Georgia 30302-4018

Users of this thesis who are not enrolled as students at Georgia State University are required to attest acceptance of the preceding stipulation by signing below. Libraries borrowing this thesis for the use of their patrons are required to see that each user records here the information requested.

NAME OF USER	ADDRESS	DATE	TYPE OF USE (EXAMINATION ONLY OR COPYING)
			,

#### VITA

Name: Anthony C. Gregory II

**Address:** 6871 Peachtree Dunwoody Rd. NE, Apt 210

Atlanta, GA, 30328

#### **Education:**

2006 Bachelor of Science, Biology, Concentrations in Molecular Biology and Microbiology, Georgia State University

**Honors:** National Honor Society; Golden Key Society. Dean's List 2001-2002, 2003-2004. William G. Nolan Memorial Scholarship,

#### **Professional Experience:**

#### -Masters in Public Health

Currently in a Masters in Public Health Program at Georgia State University

#### -Linguistic

- Worked as a Spanish Medical Interpreter Volunteer at Grady Memorial Hospital for 2 years
- Fluent in Spanish, native English speaker, also speaks some French

#### -Laboratory

• I have conducted research, for 2 years, to categorize the Cytotoxic-T Lymphocyte responses to Herpes viruses.

#### -Leadership

 Worked for the US Army reserves for 7 years as a Psychological Operations Specialist which specialized in all forms of communication

#### **EMPLOYMENT HISTORY**

Masters in Public Health Graduate Research Assistant	1/07-Present
Center for Disease Control and Prevention (CDC/ATSDR) intern	3/07-6/07
Spanish Translator Volunteer, Grady Memorial Hospital	5/05-5/07
B Virus Laboratory	5/05-5/07
US Army Reserves	8/01-5/05
US Army Active	7/97-7/0

#### Table of Contents

Chapter	Page
ACKNOWLEDGEMENTS	. iv
List of Tables and Figures	x
List of Acronyms and Abbreviations	. xii
I. INTRODUCTION	. 1
Background	. 1
The Epidemiology of MS	. 2
Research Questions	3
II Literature Review	4
Overview of Factors	. 4
Genetic Factors	. 5
Biological Factors	7
Viruses as Biological Exposure Agents	8
Sunlight and Vitamin D as Physical and Chemical Agents	11
Other Environmental Factors	13
III. METHODOLOGY	16
Study Purpose	16
Data Sources	16
Data Management	. 18
Statistical Analysis	. 20
IV. RESULTS	. 22
V. DISCUSSION AND CONCLUSIONS	90
Summary of Figures and Tables	. 90
Discussion	. 91

Advantages of This Study's Methods	. 93
Limitations of This Study's Methods	94
Conclusions and Recommendations for Future Research and Policy	95
6. REFERENCES	98
APPENDIXES A-C	101

#### **List of Tables and Figures**

Page
Table 1. Multiple Sclerosis in GA by County: Self-Reported Cases and Prevalence Rates per 100,000 Population
Figure 1. Total MS Prevalence in GA, by County: Self-Reported Prevalence Rates, November 2006
Figure 2. MS Prevalence among Females in GA, by County: Self-Reported Prevalence Rates, November 2006
Figure 3. MS Prevalence among Males in GA, by County: Self-Reported Prevalence Rates, November 2006
Table 2: Bivariate Linear Regression for Self-Reported Prevalence and County Level Census Data
Table 3: Bivariate Linear Regression for Self-Reported Prevalence versus County Level Emissions Data
Table 4: Bivariate Linear Regression: Total Self-Reported Prevalence versus Criteria Air Pollutant Data
Table 5: Multivariate Linear Regression, Total MS Self-Reported Prevalence Rates and Census and NEI air pollutants
Table 6: Multivariate Linear Regression, Female MS Self-Reported Prevalence Rates and Census and NEI air pollutants
Table 7: Multivariate Linear Regression Data, Male MS Prevalence Rates and Census and NEI air pollutants
Table 8: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: Census Data and Total Self-Reported MS Cases
Table 9: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: Census Data and Female Self-Reported MS Cases
Table 10: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: Census Data and Male Self-Reported MS Cases
Table 11: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: County Level Emissions and Total MS Self-Reported Cases

Chapter	age
Table 12: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: County Level Emissions and Female Self-Reported MS Cases	
Table 13: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: County Level Emissions and Male MS Self-Reported Cases	
Table 14: Odds Ratios(OR) and 95% Confidence Interval (CI) Organized by Quart Criteria Air Pollutants and Total MS Self-Reported Cases	
Table 15: Odds Ratios(OR) and 95% Confidence Interval (CI) Organized by Quart Criteria Air Pollutants and Female MS Self-Reported Cases	
Table 16: Odds Ratios(OR) and 95% Confidence Interval (CI) Organized by Quart Criteria Air Pollutants and Male MS Self-Reported Cases	

#### List of Acronyms and Abbreviations

CI: Confidence Interval

CO: Carbon Monoxide

EAE: Experimental Autoimmune Encephalomyelitis (the mouse Multiple Sclerosis

model)

EBNA: Epstein-Barr Nuclear Antigen

EBV: Epstein-Barr Virus

EPA: United States (U.S.) Environmental Protection Agency

GA: Georgia

HLA: Human Leukocyte Antigen

HHV: Human Herpes Virus

IRB: Institutional Review Board

IgE: Immunoglobulin Type E

MS: Multiple Sclerosis

NMSS-GA: Multiple Sclerosis Society- Georgia Chapter

NEI: National Emissions Inventory (of the U.S. EPA)

NO<sub>x</sub>: Nitrogen Oxides

OR: Odds Ratio (*Please note, in this study, OR signified Prevalence Odds Ratio*)

PM: Particulate Matter

RR: Relative Risk

RRMS: Relapsing Remitting Multiple Sclerosis

PPMS: Primary Progressive Multiple Sclerosis

SO<sub>2</sub>: Sulfur Dioxide

TRI: Toxic Release Inventory (of the U.S. EPA)

**VOC: Volatile Organic Compounds** 

#### I. Introduction

Background

Throughout time, physicians and researchers have studied thousands of diseases and conditions. In that process, they have found many causes, influencing factors and treatments. However, some diseases still present challenges with respect to determining their causes. One of those challenges is Multiple Sclerosis (MS), a disease that was first identified hundreds of years ago and that affects millions of people across the globe. MS is the most common demyelinating disease of the central nervous system. In 1992, Anderson et al (Anderson et al. 1992) estimated in the United States between 250,000 and 350,000 people have been affected by this disease.

MS is a disease in which the body's immune cells attack neuronal myelin sheaths. (Keegan and Noseworthy 2002) In healthy individuals, these sheaths help the neurons to conduct messages to other neurons by covering most of the outside of the neuron's axon. This covering mediates faster neurological signals. Due to continued attacks in MS, the myelin is destroyed, and, hence, the nerves are less efficient in conducting electrical signals. Local inflammation may further damage the neuronal axons, the long extensions that help conduct electrical signals from one part of the neuron to the other. The continued degeneration of the myelin may lead to extreme fatigue and progressively less ability to move. This process is suspected to be mediated largely by T-helper-1 cells (Th1). The degeneration seen in this disease is

primarily diagnosed by magnetic resonance imaging (MRI); however, MRI specificity makes diagnosis difficult. (Fazekas et al. 1999)

#### The Epidemiology of MS

The epidemiology of MS is also important to consider. In areas with higher MS prevalence, like Europe and North America, the prevalence of MS seems to vary from 60 to 200 per 100,000 population, while in lower MS areas, like Japan, these prevalence rates are around 6-20 per 100,000. Of note, even though the prevalence of MS is lower in Asian countries, Japan has the highest prevalence of MS among Asian countries. Japan is also one of the most developed countries in Asia. Thus, there are geographical differences in the MS distribution. These differences are probably due to a variety of genetic and environmental characteristics of these areas. (Sospedra and Martin 2005; Sospedra et al. 2005)

A descriptive MS study was conducted by Williamson et al. (Williamson 2007) in 19 counties surrounding Lubbock, Texas. This study found the prevalence of MS in this region per 100,000 population was 42.8 overall, 68.6 in females and 16.6 in males. Also, the prevalence per 100,000 population was 56.0 in non-Hispanic whites, 22.1 in non-Hispanic blacks, and 11.2 in Hispanics. They also reported the 40-59 year-old age groups had the highest prevalence of MS; the 40-49 year-old group had 103.1 per 100,000 and the 50-59 year-old group had 119.9 per 100,000. (Williamson et al 2007) In addition to this, the prevalence of MS has varied widely

throughout the world (Baum and Rothschild 1981; Mayer 1981) (Rosati 2001) (de la Maza et al. 2000), (Kurtzke 2000)

There seems to be little definitive information as to the etiology of MS. Although there are many theories about the exact role of genetics, immune cells, viruses, and other possible agents, there is little consensus about a single casual element for MS. Thus, it is evident MS is a complex disease, and MS embodies a multitude of genetic and environmental factors like exposure to certain viruses, chemicals, and possibly environmental air pollutants subject to regulation.

#### Research Questions

The motivation for this study comes from the knowledge of the lack of geographical models of MS and the lack of information as to the potential impacts of environmental air pollutants in MS. With these focuses in mind, several research questions were created for this thesis.

- 1. What is the distribution of the prevalence of MS in GA?
- 2. How does this distribution relate to census data?
- 3. How does the distribution of MS relate to environmental air pollutants?
- 4. Using both environmental air pollutants and census data, what is the best predictive model for the MS distribution in GA?
- 5. Is there a dose-response-like relationship between the MS distribution in GA and specific, census selected data characteristics or environmental air pollutants?

#### II. Literature Review

Overview of Factors

Many factors, both genetic and environmental, have been associated with MS (Hogancamp et al. 1997). Of the genetic factors, twin studies have been instrumental. These studies have compared monozygotic twins (or identical twins, which have identical genetic information) and dizygotic twins (or paternal or non-identical twins, which have different genetic information) to determine if there is a significant genetic role for MS. Gender-related genetic factors have also been identified.

Aside from the genetic role, many biological factors have also been identified through scientific research. This research is commonly done in animal models for MS as a surrogate to conducting these studies in humans.

Environmental factors have also shown significant association with MS. For example, Mayer et al. (1981) found mean annual hours of sunshine and mean annual temperature were protective of MS. Conversely, infection with certain viruses (Keegan et al. 2002, Berti et al. 2002, (Soldan et al. 1997), distance from the equator (Mayer 1981), low average amount of daily sunlight (Freedman et al. 2000), lack of Vitamin D (Cantorna et al. 1996; Cantorna et al. 1998; Spach and Hayes 2005), exposure to heavy metals (Henry et al. 2007), per capita steel consumption (Mayer 1981), number of physicians per capita (Mayer 1981), literacy rate (Mayer 1981), gross energy consumption, gross national product, per capita and food consumption are associated with higher rates of MS (Mayer 1981). Among the many factors that may play a role in MS, viruses have been commonly cited as a common factor that

may influence the acquisition and progression of MS. More details by category now follow.

#### Genetic Factors

There have been several genetic clues to the etiology of MS. Twin studies have been an important part in the etiology of MS, because identical twins have identical genetic information, while fraternal twins do not. Twins are both the same age, and, assuming that they have similar environmental exposures, one can estimate the relative role of genetics in a disease. In twin studies conducted for MS, a significantly higher prevalence of MS was found in monozygotic twins when compared to dizygotic twins (de Rezende and Arruda 1996; Hansen et al. 2005; Keegan and Noseworthy 2002). First-degree relatives of affected people had an about 20 to 50-fold (2%–5%) higher risk to develop MS. Also, concordance rates in monozygotic twins varied between 20% and 35% in different studies, and recent studies estimated the concordance rate at 25%. (Dyment et al. 2004) (Ebers et al. 1996) This higher prevalence of MS in monozygotic twins suggested there was a significant role of genetics in MS.

The population of people affected by MS has been categorized. Females are more susceptible than males (1.6:1), but this difference is evident in Relapsing Remitting Multiple Sclerosis (RRMS), in which the person recovers completely, but not in Primary Progressive Multiple Sclerosis (PPMS), in which the patient never completely recovers. (Keegan and Noseworthy 2002) (Thompson et al. 1997) (Weinshenker 1998)

Aside from the higher risk of MS in women, other studies have supported the notion female hormones may play a role in MS. In one of these studies, Runmarker and Anderson (1995) analyzed the risk of MS before, during and after pregnancy in a cohort of Swedish women. They found there was a significantly decreased risk of MS during pregnancy. Furthermore, they found after birth the risk did not differ significantly from periods when the women were not pregnant. Also, they found women who have given birth had a lower risk of MS than women who had not given birth. Lastly, they found a significantly lower risk of progressing from RRMS to PPMS in the group of women who became pregnant after the onset of MS, when compared to non-pregnant control patients.

These findings suggested there may be some role of pregnancy hormones in MS, and other studies have also reported this same relationship. Sicotte et al. (2002) found treatment of RRMS patients with estradiol (a pregnancy hormone) resulted in significant decreases in the number of lesions found during monthly magnetic resonance imaging. This same protective effect of estradiol was seen in a later experiment conducted by Soldan et al. (Soldan 2003). Both studies surmised this change may be due to the hormone's effect of changing T-helper 1 immune responses to T-helper-2 immune responses. (NL Sicotte 2002; Soldan 2003)

In addition to gender-roles, some racial and ethnic groups have been associated with generally higher rates of MS. According to Rosati et al. (2001), MS was rare among Samis, Turkmen, Uzbeks, Kazakhs, Kirghizis, native Siberians, North and South Amerindians, Chinese, Japanese, African blacks and New Zealand

Maoris. Contrarily, they also saw a higher risk for MS among Sardinians, Parsis and Palestinians. Also, other researchers have reported a significantly higher risk for MS in people of Northern European descent, although there was a lower prevalence of MS in New Zealand (Keegan and Noseworthy 2002).

#### Biological Factors

Scientific experiments have been used to attempt to discover more biological clues to the cause of MS. The animal-model of MS is known as Experimental Autoimmune Encephalomyelitis (EAE). To induce EAE in mice, scientists exposed them to myelin basic protein, a protein similar to proteins found in myelin of most animals. Thus, the immune cells will react to this protein and also to natural myelin surrounding murine neurons. Catorna et al. (1996, 1998) and Spach et al. (2005) conducted experiments in the mouse model for Multiple Sclerosis.

EAE studies have been largely a part of determining the immunological basis for MS. Through EAE studies, scientists determined that CD4+ T-helper 1 cells might be mediating this disease. This determination was made upon realizing that transferring these CD4 Th1 cells could transfer EAE from an affected mouse to an unaffected mouse. Transferring antibodies has not been effective for transferring EAE, and transferring CD8+ cells has only been effective in transferring EAE in two instances. The genetic sequence of human leukocyte antigen (HLA) class II molecules is the strongest genetic risk for MS, and some studies also supported the idea that CD4 Th1 mediates MS. (Sospedra and Martin 2005; Sospedra et al. 2005) (Oksenberg et al. 2001) Furthermore, there may be variation in these HLA genes, as

Ito et al. (Ito et al. 1998) found increased susceptibility with the western-type MS but not with the Japanese-type MS.

Viruses as Biological Exposure Agents

In addition to the EAE animal model, there is Theiler's murine encephalomyelitis virus model, in which a virus is deposited into susceptible mice, the mice develops an encephalomyelitis, and this leads to neuron and myelin degeneration (Dal Canto et al. 1995).

In addition to the virus-mediated animal MS model, several authors have proposed viruses may cause or exacerbate MS. Soldan and others (1997) showed that IgM, a common antibody for early stages of infection for Human Herpes virus 6 (HHV-6), was increased in RRMS when compared to PPMS. They also showed HHV-6 serum DNA in MS patients (Soldan et al. 1997). Berti et al. (2002) confirmed the presence of HHV-6 DNA in MS patients, and they stated the presence of HHV-6 DNA was correlated with MS progression. These studies, together, suggested there may be a role of HHV-6 in MS. In addition, the association between the viral DNA titers and MS progression suggested as the amount of total virus burden increased, there may also be increased severity of MS. (Berti et al. 2002)

Several other studies also lend credibility to the role of HHV-6 in the etiology of MS (Akhyani et al. 2000; Kong et al. 2003; Tejada-Simon et al. 2003; Virtanen et al. 2005). In a previous study, Berti et al. (2000) found an increased frequency of HHV-6 DNA in the serum, saliva, urine, and peripheral blood lymphocytes when compared to controls. Kong et al. (2003) found HHV-6 may play a role in

degenerating oligodendrocytes, the cells that help to nurture and create myelin. Tejad-Simpson et al. (2003) found 50% of T-cells were treated with HHV-6 residues cross-reacted with myelin basic protein. This suggested T-cells that have seen foreign HHV-6 proteins may have mistakenly learned that myelin was also foreign; therefore, these T-cells may have mounted an immune response to neuronal myelin. In a study about MS and control brain tissue, Virtanen et al. (2005) found HHV-6 in the same place as MS lesions more often then in control brains.

One study found HHV-6 was not related to MS. In this study, Tuke et al. (2004) found there was no difference in distribution, virus type, or quantity of HHV-6 in MS patients when compared with control patients.

In addition to the possible role of HHV-6, Epstein Barr Virus (EBV) has also been associated with MS. (Hernan et al. 2001; Levin et al. 2003) They determined the presence of a Epstein Barr Virus Nuclear Antigen (EBNA) was associated with an increased risk of MS (Hernan et al. 2001). In a similar study, Levin et al. (2003) also found the risk of MS increased as the antibody titer to EBV antigens increased. When these authors compared the lowest EBNA titer level with the highest, they found the relative risk for MS was 33.9 (95% CI=4.1-283) (Levin et al. 2003; Levin et al. 2005). Another researcher, Lunemann et al. (2006), found EBNA-specific CD4+ memory T Cells (T-helper cells) were much more common than any other type of EBV antigen in MS patients who were seropositive for EBV.

Several other authors have presented studies supporting the role of EBV in the etiology of MS. (Cepok et al. 2005; Haahr et al. 2004; Lunemann et al. 2006; Sumaya

et al. 1985; Wandinger et al. 2000) Sumaya et al. (1985) found a greater antibody response to EBV antigen in MS patients when compared to controls. Furthermore, Wandinger et al. (2000) found antibodies to EBV in 100% of all the MS patients in their study. Cepok et al. (2005) found immune-cell reactivity for EBNA was much higher in the serum and cerebrospinal fluid of MS patients when compared with control patients. Haahr et al. (2004) found a high-MS prevalence area had more than one third of children who were seronegative for EBV. They also found 100% of MS patients in their sample were positive for EBV. Thus, they posited EBV was transmitted to a large portion of the population during or after puberty in the high-MS prevalence area. Finally, Wagner et al. (2004) found there was an increased risk of MS in people who have EBV in the serum, even after adjusting or smoking, ancestry, and latitude of residence.

There seems to be some similarities between both HHV-6 and EBV. Both viruses are herpes viruses, and they both replicate slowly. Beta herpes viruses, like HHV-6, are known for infecting cells, and cells infected typically become larger. EBV, also known as Human Herpes Virus 4, is a gamma herpes virus, which is known for being associated with lymphocytes. Also, gamma herpes viruses were found to be able to induce cell-proliferation. This is important because T-cell proliferation and activation usually precedes T-cell immune responses. Also, proliferation and activation of other cells may lead to the characteristic inflammation found in many MS patients. (Hong et al. 2000; Levin et al. 2005)

Sunlight and Vitamin D as Physical and Chemical Agents

Other environmental exposures like mean hours of sunshine and vitamin D have been associated with a lower prevalence of MS (Mayer 1981; Munger et al. 2004) Mayer et al. (1981) found mean annual hours of sunshine and mean annual temperature. Freedman et al. studied the deaths from MS for the years of 1984 and 1995 in 24 states of the U.S. (2000), and found a 43% decrease in MS rates amongst those with the highest category of sun exposure. Through their analysis, they also found under high sunlight exposures black and white men and women received a protective effect of sunlight. This information suggested sunlight had a significant physiological role that helped to mitigate MS in these two racial groups.

Several authors supported the potential role of average incident sunlight in MS by showing a gradually increasing diagnosis of MS in the higher latitudes (Keegan et al. 2002, Mayer et al. 1981, Rosati et al. 2001). Seasonal fluctuations may influence MS, with most attacks occurring in the most extreme temperatures of summer and winter (Ogawa et al. 2004). Taken together, these authors suggested some environmental factors related to sunlight may affect MS prevalence.

To characterize the relationship between sunlight and the incidence of MS, Munger et al. (2004) studied the role of vitamin D, which is primarily produced by the body through sunlight exposure or ingested in the form of daily vitamins.

Through the Nurses Health Study I and II, they analyzed the role of behaviors, like taking daily vitamins, on the incidence of several diseases. They found there was a significantly lower relative risk (RR=0.59; 95% CI=0.38 to 0.91; p for trend = 0.006)

for MS in women who are taking 400 IU of vitamin D per day, when compared to those who did not take this supplement. They also found this relationship was not seen for vitamin D taken through food, and this relationship was dose dependent. These analyses suggested the ingested daily vitamin may have protective effects for MS. The dose response suggested environmental exposure of sunlight and the dose of vitamin D both have effects on the incidence and prevalence of MS.

Cantorna et al (1996) found 20 ng/day of Vitamin D completely prevented disability in EAE induced mice; the mice were given a calcium diet, but only the experimental mice were given 20 ng of vitamin D. Also, this regimen seemed to halt progression of EAE after it reached the earliest stage (limp tail in mice), and when the vitamin D was removed, EAE progression returned to the normal level. Cantorna et al (1998) further categorized the role of vitamin D in a later experiment, where they reported systemic and local increases in two anti-inflammatory cytokines, facilitated by Vitamin D, may be responsible for blocking EAE. These experiments suggested that vitamin D may not only prevent Multiple Sclerosis in some cases, but also vitamin D may be able to halt or slow the progression of MS in mice. Although we are unsure if this experiment has the same effects in humans, it provides great insight into the interaction between vitamin D and MS.

In a separate EAE experiment, Spach et al. (2005) found ultraviolet-light exposed, vitamin D(3)-fed female mice had less clinical, histopathological and immunological signs of EAE than ovariectomized females or intact or castrated males. They summarized this effect seemed to show synergy between vitamin D and

ovarian tissue with regard to EAE inhibition. They further hypothesized an ovarian hormone inhibited CYP24A1 expression, which encodes a vitamin D inactivating enzyme, in the spinal cord, and so 1,25-dihydroxyvitamin D (vitamin D) accumulated and resolved the inflammation before severe EAE developed. Lastly, they speculated sunlight deprivation would increase the MS risk more significantly in women than in men, which may contribute to the unexplained higher MS incidence in women than in men. (Spach and Hayes 2005) If this study can be extrapolated to humans, it may suggest women may benefit more than men from taking vitamin D, and it may help to partially explain the higher incidence of MS in women.

#### Other Environmental Factors

Several other types of environmental exposures have been shown to be potentially associated with the prevalence and incidence rates of MS. Seasonal fluctuations have been associated with MS in a study conducted in Japan by Ogawa et al. (2004). They found optico-spinal and brain MS complications were significantly higher in Japan during the coldest (January and February) and hottest months (July and August) of the year. They also theorized common infections occurring during the cold season, such as sinusitis, may play a role in these seasonal fluctuations of MS. (Ogawa et al. 2004)

In another study, Edwards et al. (1998) suggested there was a relationship between infections and MS exacerbations. They reported a twofold increase in MS exacerbations (p=0.004, OR 2.0, CI 1.3–3.2) during symptomatic upper respiratory track infections that lasted more than 24 hours when compared to periods without

symptomatic upper respiratory tract infections. Thus, these researchers found upper respiratory tract infections, which may be caused by viruses, may also affect the frequency of MS symptoms.

Henry et al. (2007) analyzed MS in cohorts based in two Texas schools. They reported 22 of the women in their study had MS, and they found a crude prevalence estimate of 411 per 100,000 population. Also, they established three of the 22 women, who had childhood hair samples collected and archived to test for heavy metals exposure, had elevated levels of lead and mercury in their childhood hair. Several authors have also proposed a potential role of heavy metals in new clusters of MS cases (Ingalls 1986) (Landrigan et al. 1975). In addition to this, Williamson and Henry (2004) acknowledged there has been community concern over the possible role of toxins from hazardous waste sites in new cases of MS.

Few studies have focused on the roles of environmental air in the incidence and progression of MS. One study on this subject was conducted in Finland by Oikonen et al. (2003). In this study, the role of ambient air quality was found to be associated with MS prevalence. In this study, the researchers found MS relapse was associated with PM-10 concentrations in southern Finland. They found there was a strong interaction between PM-10, CO and acidic gases (NO<sub>x</sub> and SO<sub>2</sub>). Thus, they used multivariate logistic regression to attempt to differentiate the roles of acidic gases and the roles of PM-10 from each other. They found the odds of relapse was 4.143 (p<0.001) following a one-month lag after peak levels of PM-10. They also found an even larger odds ratio for acidic gases one month after peak levels

(OR=9.285, 95% CI=2.744, 31.418). These results, however, must be interpreted with caution due to small sample sizes and crude exposure estimates.

Also, another study found that environmental exposure to radon may be a contributory factor in MS. In this study, Bolviken et al (Bolviken et al. 2003) assessed the correlation between radon exposure and MS, and they found a positive correlation (p< 0.01). They surmised that Radon inhaled in air maybe contributory factor to MS.

These data suggested that MS relapse may be associated with high rates of environmental air pollutants in Finland. With this in mind, the purpose of this study will be to assess the suspected role of environmental air pollutants in MS, using data on source emissions and MS in the state of Georgia by county.

#### III. Methods

The GSU Institutional Review Board (IRB) approved this study in February 2007 (approved protocol # H07341). To receive approval, the methodology and supporting documents were submitted to GSU through an on-line system, IRBWise.

#### Study purpose

This study is designed to estimate the county-level prevalence and distribution of MS throughout Georgia (GA). In addition, this study will determine if the distribution of MS in GA is associated with the distribution of census data, toxic release inventory (TRI) data and source emission to outdoor air from the National Emissions Inventory database (NET) of criteria air pollutants. Also, this study will attempt to determine which of these three data relate best to the distribution of MS in GA. Lastly, this study will endeavor to determine if there is a dose response relationship between the distribution of MS and the census or air pollutant data.

#### Data Sources

The data for this study comes from the National Multiple Sclerosis Society (NMSS), the United States Census Bureau, and Scorecard.org, by Environmental Defense, which assimilated county-level air and water emissions data in the United States. (US Census 2005, Scorecard 2005) The final population for this study was 9,072,576 people, including 6247 self-reported MS cases from the NMSS-GA chapter.

MS data were received from the National Multiples Sclerosis Society's GA chapter in November 2006. These data included the self-reported MS cases of the current MS Society member's list as of November 2006. The data obtained for this study includes the number of people diagnosed with multiple sclerosis, their gender, and their county of residence. The NMSS, as part of their membership questionnaire, originally received this information.

Selected census data were retrieved for each of the 159 Georgia counties. These data included the total county population, the percent female in 2005, the 2005 female and male population estimates, the percent white and the percent black. In addition, the percent living in the same house in 1995 and 2000, the per capita income, in 1999, the square miles land area in 2000, and the people per square mile in 2005 was also obtained for this study.

Also, using scorecard.org, which provided a database of Toxic Release

Inventory (TRI) source emissions from the U.S. Environmental Protection Agency
(EPA) for various categories of toxicants and criteria air pollutants, county level air
pollutant emissions were obtained. The data used for this study included 2002 source
emissions in pounds: recognized carcinogens, suspected carcinogens, suspected
cardiovascular or blood toxicants, recognized developmental toxicants, suspected
developmental toxicants, suspected endocrine toxicants, suspected endocrine
toxicants, suspected immunotoxicants, suspected kidney toxicants, suspected
gastrointestinal or liver toxicants, suspected muscoskeletal toxicants, suspected

neurotoxicants, recognized reproductive toxicants, suspected reproductive toxicants suspected respiratory toxicants, suspected skin and sense toxicants, acetaldehyde, ammonia, formaldehyde, manganese, and lead and lead compounds.

In addition to these, the National Emissions Inventory (NEI) Air Pollutant Emissions Trends data, which were measurements led by the U.S. EPA, were obtained through scorecard.org. NEI is a collection of source emissions to outdoor air from state, local, tribal and industrial measures. From NEI, this study used criteria air pollutants reported for 1999 in tons: carbon monoxide, nitrogen oxides, particulate matter (PM) 2.5, PM 10, sulfur dioxide, and volatile organic compounds (total, selected individual). Each compound was surveyed for mobile, area, and point sources; the three source categories together were termed "all sources."

It is important to state this study did not assume source emissions are the same as measured concentrations in outdoor air or actual human exposure. Rather, this study used source emissions as a surrogate measure for exposure to environmental air pollution, selected groups of toxicants and criteria air pollutants subject to regulation.

#### Data Management

For this study, a MS case was defined as a person being diagnosed previously with MS, according to the NMSS entrance survey, and residing in GA. Furthermore, the number of controls was calculated using the 2005 U.S. Census population for each county, minus the number of MS cases in that county. Lastly, prevalence signified the county self-reported case prevalence per 100,000 population.

Several assumptions were necessary for this analysis. We assumed the NMSS-GA member list included accurate data members answered truthfully as to their diagnosis of MS and their gender. Also, we assumed the TRI and NEI data were received and displayed in scorecard.org's website and it was accurately reported by the private and public sector per the Emergency Planning and Community Right-to-Know Act of 1986, which established TRI.

This study used cross-sectional data at several different time points. The study was designed to determine the prevalence of MS in GA and to explore the relationship between environmental air pollutants and MS in GA. With that purpose in mind, we used November 2006 cross sectional data from the MS Society's GA chapter. These data were compared to the county-level emissions data using both regression (a measurement of association) and odds ratios.

Female and male prevalence rates were calculated for each of the counties in GA by dividing the number of cases in each county by the total county population and multiplying this number by 100,000. Thus, each of the prevalence ratios shows a population-density-controlled rate for each county. The database was stored in both SPSS version 14 (SPSS 2005) and Microsoft Excel 2004 (Microsoft-Corporation 2004). These prevalence rates were made into a geographical image using CorelDraw by tracing the image of a map of GA and placing the prevalence rates in their respective counties. After this was done, each county was shaded according to

the county's prevalence rate. This same process was done for total, female, and male, MS rates in GA per 100,000 population.

#### Statistical Analysis

Statistical analysis was done using both SPSS version 14 (2005) and Microsoft Excel 2004 (Microsoft-Corporation 2004). Bivariate linear regression was used to determine association between the prevalence rates and each of the census data or emissions data variables. Standardized beta values were calculated for each of these associations, and the statistical significance was calculated using a two-tailed t-test. Finally, R-squared values were also calculated for each model. R-squared represents the amount of variance in the model (this model's female self reported MS prevalence in GA) that can be attributed to the study characteristic (eg. Percent female 2005).

Stepwise multivariate linear regression was calculated using SPSS. This was done with an entry criterion of a p-value of 0.05 and an exit criterion of a p-value of 0.10. With this method, we analyzed which of the census and NEI variables were most related to the county MS prevalence in GA, overall (total) and by gender.

Odds ratios (ORs) were calculated for each of the census characteristics, each of the toxicants, and each of the criteria air pollutants. It is important to mention the odds ratios in this study referred to the prevalence odds ratio because the cases used in this study indicated prevalence, not incidence. The counties with data for the variable being analyzed were separated into quartiles by SPSS for Windows version

14. The total number of cases or controls was added for each of the quartiles and these numbers were used to calculate the ORs. The lowest quartile was always used as the reference group. Then, the 2<sup>nd</sup>, 3<sup>rd</sup> and 4<sup>th</sup> quartiles were compared to the lowest quartile using EPI Info version 6.04 (CDC 2001). Finally, data were analyzed statistically at the 0.05, 0.01, 0.001, and 0.0001 levels. Once all of the analyses were done, the tables were constructed using Microsoft Excel 2004.

#### **IV.** Results

Table 1 shows the Multiple Sclerosis cases and self-reported-case prevalence rates by county. In the 159 counties in GA, the prevalence rates varied widely; the rates were not equally distributed from county to county. Also, these self-reported cases are 71.5% female, 22.3% male, and 6.2% unknown gender.

Table 4 Multiple Co	dorocio in	CA by Co	untur Colf I	Departed Case	a and Dravalan	aa Dataa nar 100	000 Deputation
Table 1. Multiple Sc	Nov.	Nov.	Nov.	Nov. 2005	Nov. 2005	ce Rates per 100 Nov. 2005	Nov. 2005
	2005	2005	2005	unknown	total	female	male
	total	female	male	gender	prevalence	prevalence	prevalence
	cases	cases	cases	cases	rates* (per	rates* (per	rates* (per
	(N=	(N=	(N=	(N=389)	100,000	100,000	100,000 men)
County name	6247)	4466)	1392)		people)	women)	
Appling	3	3	0	0	17	33	0
Atkinson	2	1	1	0	25	25	25
Bacon	1	1	0	0	10	19	0
Baker	0	0	0	0	0	0	0
Baldwin	18	8	9	1	40	38	37
Banks	8	6	2	0	50	76	25
Barrow	41	32	8	1	68	107	27
Bartow	60	44	13	3	67	98	29
Ben Hill	7	4	3	0	40	44	36
Berrien	9	7	2	0	54	83	24

Table 1. Multiple S	clerosis in	GA by Cou	unty: Self-I	Reported Case	s and Prevalen	ce Rates per 100	),000 Population.
•	Nov.	Nov.	Nov.	Nov. 2005	Nov. 2005	Nov. 2005	Nov. 2005
	2005	2005	2005	unknown	total	female	male
	total	female	male	gender	prevalence	prevalence	prevalence
	cases	cases	cases	cases	rates* (per	rates* (per	rates* (per
	(N=	(N=	(N=	(N=389)	100,000	100,000	100,000 men)
County name	6247)	4466)	1392)		people)	women)	,
Bibb	110	74	28	8	71	89	39
Bleckley	3	2	0	1	25	32	0
Brantley	5	3	1	1	32	39	13
Brooks	4	4	0	0	24	47	0
Bryan	23	17	3	3	81	118	21
Bulloch	26	21	2	3	42	67	7
Burke	5	4	0	1	21	33	0
Butts	17	10	5	2	81	102	45
Calhoun	3	3	0	0	50	117	0
Camden	40	33	5	2	87	144	22
Candler	7	3	4	0	68	58	77
Carroll	78	62	13	3	74	116	25
Catoosa	0	0	0	0	0	0	0
Charlton	9	7	1	1	83	139	17
Chatham	160	132	26	2	67	107	23
Chattahoochee	2	2	0	0	14	32	0
Chattooga	13	11	1	1	49	86	7
Cherokee	179	128	40	11	97	140	43
Clarke	68	46	10	12	65	87	19
Clay	0	0	0	0	0	0	0
Clayton	129	91	32	6	48	66	24

Table 1. Multiple Sc	elerosis in	GA by Cou	ınty: Self-F	Reported Case	s and Prevalen	ce Rates per 100	,000 Population.
	Nov.	Nov.	Nov.	Nov. 2005	Nov. 2005	Nov. 2005	Nov. 2005
	2005	2005	2005	unknown	total	female	male
	total	female	male	gender	prevalence	prevalence	prevalence
	cases	cases	cases	cases	rates* (per	rates* (per	rates* (per
	(N=	(N=	(N=	(N=389)	100,000	100,000	100,000 men)
County name	6247)	4466)	1392)		people)	women)	
Clinch	5	2	3	0	71	57	87
Cobb	670	473	154	43	101	142	47
Coffee	8	6	2	0	20	30	10
Colquitt	26	17	6	3	59	78	27
Columbia	95	67	21	7	92	126	41
Cook	6	6	0	0	37	71	0
Coweta	92	70	12	10	84	127	22
Crawford	6	3	2	1	47	47	31
Crisp	8	6	2	0	36	52	19
Dade	0	0	0	0	0	0	0
Dawson	18	12	3	3	91	123	30
Decatur	8	7	1	0	28	47	7
Dekalb	547	373	138	36	81	107	42
Dodge	9	7	1	1	46	75	10
Dooly	3	1	2	0	26	18	33
Dougherty	44	38	5	1	46	75	11
Douglas	92	64	26	2	82	112	47
Early	5	5	0	0	41	78	0
Echols	1	0	0	1	24	0	0
Effingham	29	24	4	1	62	102	17
Elbert	9	8	1	0	43	74	10

Table 1. Multiple Sclerosis in GA by County: Self-Reported Cases and Prevalence Rates per 100,000 Population.										
Table 1. Multiple So	Nov.	Nov.	Nov.	Reported Case Nov. 2005	Nov. 2005	Nov. 2005	Nov. 2005			
	2005	2005	2005	unknown	total	female	male			
	total	female	male			prevalence				
				gender	prevalence rates* (per	rates* (per	prevalence rates* (per			
	cases (N=	cases (N=	cases (N=	Cases	100,000	100,000	100,000 men)			
County name	`	`	`	(N=389)	· ·	•	100,000 mem)			
County name Emanuel	6247)	4466)	1392)	1	people) 41	women)	28			
	9	5 1	3	0		44 17	54			
Evans	21		2		35					
Fannin		19		0	96	169	19			
Fayette	123	84	30	9	118	158	59			
Floyd	72	54	16	2	76	112	35			
Forsyth	144	96	36	12	103	139	50			
Franklin	16	14	2	0	74	126	19			
Fulton	745	511	177	57	81	111	39			
Gilmer	11	9	2	0	40	67	14			
Glascock	3	3	0	0	111	218	0			
Glynn	50	32	12	6	70	86	35			
Gordon	29	19	9	1	58	76	36			
Grady	11	8	2	1	45	63	17			
Greene	15	12	2	1	96	148	26			
Gwinnett	531	382	120	29	73	107	33			
Habersham	17	14	3	0	43	73	15			
Hall	87	66	17	4	52	82	20			
Hancock	3	3	0	0	31	67	0			
Haralson	21	21	0	0	74	146	0			
Harris	15	7	6	2	54	50	44			
Hart	8	7	1	0	33	57	8			

Table 1. Multiple Sclerosis in GA by County: Self-Reported Cases and Prevalence Rates per 100,000 Population.											
Table 1. Multiple Sc	1										
	Nov.	Nov.	Nov.	Nov. 2005	Nov. 2005	Nov. 2005	Nov. 2005				
	2005	2005	2005	unknown	total	female	male				
	total	female	male	gender	prevalence	prevalence	prevalence				
	cases	cases	cases	cases	rates* (per	rates* (per	rates* (per				
County name	(N=	(N=	(N=	(N=389)	100,000	100,000	100,000 men)				
County name	6247)	4466)	1392)	^	people)	women)	10				
Heard	6	5	1	0	53	87	18				
Henry	142	105	27	10	85	124	32				
Houston	98	74	18	6	78	116	29				
Irwin	2	2	0	0	20	39	0				
Jackson	54	34	17	3	103	131	65				
Jasper	7	5	1	1	53	75	15				
Jeff Davis	3	1	0	2	23	15	0				
Jefferson	8	4	1	3	47	45	13				
Jenkins	3	3	0	0	34	67	0				
Johnson	1	11	0	0	10	23	0				
Jones	25	15	9	1	93	110	68				
Lamar	16	12	2	2	98	142	25				
Lanier	3	3	0	0	40	81	0				
Laurens	23	17	6	0	49	70	26				
Lee	20	12	6	2	64	78	38				
Liberty	23	18	5	0	40	65	17				
Lincoln	2	1	0	1	24	24	0				
Long	1	1	0	0	9	18	0				
Lowndes	52	42	10	0	54	87	21				
Lumpkin	10	6	3	1	41	49	25				
Macon	3	3	0	0	22	43	0				

Table 1. Multiple Sclerosis in GA by County: Self-Reported Cases and Prevalence Rates per 100,000 Population.										
Table 1. Multiple 3	Nov. 2005	Nov. 2005	Nov. 2005	Nov. 2005 unknown	Nov. 2005 total	Nov. 2005 female	Nov. 2005 male			
	total	female	male	gender	prevalence	prevalence	prevalence			
	cases (N=	cases (N=	cases (N=	cases (N=389)	rates* (per 100,000	rates* (per 100,000	rates* (per 100,000 men)			
County name	6247)	(1 <b>1–</b> 4466)	1392)	(14-369)	people)	women)	100,000 mem			
Madison	25	18	6	1	92	131	44			
Marion	4	3	1	0	55	83	27			
Mcduffie	15	9	6	0	69	79	58			
Mcintosh	6	3	3	0	54	53	55			
Meriwether	16	12	4	0	70	101	36			
Miller	1	1	0	0	16	31	0			
Mitchell	8	5	3	0	34	43	25			
Monroe	11	10	1	0	46	85	8			
Montgomery	5	4	1	0	56	92	22			
Morgan	17	10	3	4	97	111	35			
Murray	0	0	0	0	0	0	0			
Muscogee	127	92	28	7	69	94	32			
Newton	57	43	10	4	66	97	23			
Oconee	30	24	4	2	101	159	27			
Oglethorpe	10	7	3	0	73	101	45			
Paulding	78	63	12	3	69	113	21			
Peach	15	9	6	0	60	71	50			
Pickens	16	10	3	3	56	69	21			
Pierce	5 10	2 8	2	1	29 62	23	24 12			
Pike Polk	20	13	3	4	49	100 65	15			
FUIK	<sub>1</sub> 20	13	3	4	49	65	15			

Table 1. Multiple Sclerosis in GA by County: Self-Reported Cases and Prevalence Rates per 100,000 Population.											
County name	Nov. 2005 total cases (N= 6247)	Nov. 2005 female cases (N= 4466)	Nov. 2005 male cases (N= 1392)	Nov. 2005 unknown gender cases (N=389)	Nov. 2005 total prevalence rates* (per 100,000 people)	Nov. 2005 female prevalence rates* (per 100,000 women)	Nov. 2005 male prevalence rates* (per 100,000 men)				
Pulaski	8	6	2	0	82	109	47				
Putnam	11	7	4	0	55	70	41				
Quitman	1	1	0	0	41	77	0				
Rabun	11	7	2	2	68	87	25				
Randolph	5	2	3	0	68	51	89				
Richmond	133	92	36	5	68	90	38				
Rockdale	62	42	16	4	79	107	41				
Schley	2	2	0	0	49	94	0				
Screven	1	1	0	0	6	13	0				
Seminole	4	2	1	1	43	41	23				
Spalding	35	27	5	3	57	86	17				
Stephens	10	7	1	2	40	54	8				
Stewart	0	0	0	0	0	0	0				
Sumter	14	10	4	0	43	58	25				
Talbot	8	6	1	1	119	170	32				
Taliaferro	0	0	0	0	0	0	0				
Tattnall	9	6	3	0	39	61	22				
Taylor	3	3	0	0	34	67	0				
Telfair	3	3	0	0	23	53	0				
Terrell	4	4	0	0	37	72	0				
Thomas	24	21	2	1	54	89	9				

Table 1. Multiple So	clerosis in	GA by Cou	ınty: Self-I	Reported Case	s and Prevalen		,000 Population.			
	Nov.	Nov.	Nov.	Nov. 2005	Nov. 2005	Nov. 2005	Nov. 2005			
	2005	2005	2005	unknown	total	female	male			
	total	female	male	gender	prevalence	prevalence	prevalence			
	cases	cases	cases	cases	rates* (per	rates* (per	rates* (per			
	(N=	(N=	(N=	(N=389)	100,000	100,000	100,000 men)			
County name	6247)	4466)	1392)		people)	women)				
Tift	15	11	2	2	37	53	10			
Toombs	11	10	0	1	40	71	0			
Towns	16	12	4	0	155	223	81			
Treutlen	3	1	2	0	44	30	58			
Troup	29	24	4	1	47	74	13			
Turner	4	3	1	0	42	61	22			
Twiggs	6	3	3	0	58	56	61			
Union	18	12	6	0	91	120	62			
Upson	23	16	5	2	83	111	38			
Walker	3	3	0	0	5	9	0			
Walton	59	41	16	2	78	106	43			
Ware	14	10	4	0	41	58	23			
Warren	3	3	0	0	49	93	0			
Washington	7	6	1	0	35	54	11			
Wayne	12	6	6	0	42	44	41			
Webster	1	1	0	0	44	88	0			
Wheeler	1	0	0	1	15	0	0			
White	17	12	3	2	71	99	25			
Whitfield	1	1	0	0	1	2	0			
Wilcox	3	2	1	0	34	53	20			
Wilkes	5	4	1	0	48	74	20			

Table 1. Multiple Sclerosis in GA by County: Self-Reported Cases and Prevalence Rates per 100,000 Population.										
·	Nov. 2005 total cases (N=	Nov. 2005 female cases (N=	Nov. 2005 male cases (N=	Nov. 2005 unknown gender cases (N=389)	Nov. 2005 total prevalence rates* (per 100,000	Nov. 2005 female prevalence rates* (per 100,000	Nov. 2005 male prevalence rates* (per 100,000 men)			
County name	6247)	4466)	1392)		people)	women)				
Wilkinson	10	5	4	1	99	94	83			
Worth	14	6	8	0	64	52	76			
Mean	39.2	28.1	8.8	2.4	53.2	75.9	23.3			
Standard Dev.	101.5	71.1	23.9	6.9	28.5	42.9	20.9			
Median	10	7	2	0	49.4	74.4	21.8			


<sup>\*</sup>Prevalence rates are calculated using the number of cases in the county divided by the county's 2005 population estimate (Census-Bureau, Jan 2007) multiplied by 100,000. The denominators for the total, female, and male rates were the total county population, the female county population, and the male population, respectively.

In figure 1, the overall self-reported-case prevalence rates at the county level were usually lower than the rates seen for females (figure 2) and higher than for male rates (figure 3). Also, the darker areas seem to center around Cobb, Fulton and Dekalb County. These counties are counties of metropolitan Atlanta, the most developed and urbanized setting in Georgia.


In figure 2, the self-reported-case prevalence rates are generally higher than those seen in figures 1 and 3, which depict overall self-reported-case prevalence rates and the male self-reported-case prevalence rates, respectively. Also, the dark area seems to center around Cobb, Fulton and Dekalb County. These counties are metropolitan counties of Atlanta, the most developed and urbanized setting in Georgia.

In figure 3 the male MS self- reported case prevalence rates are markedly lower than those seen in the female MS self-reported-case prevalence rates (figure 2). Also, while there is a slight pattern of higher rates around Atlanta, the general pattern seems more dispersed for males than for females.

<u>Figure 1:</u>
<u>Total Multiple Sclerosis Self-Reported Prevalence Rates in GA, by County: Crude Rates as of November 2006</u>



<u>Figure 2:</u>
<u>Multiple Sclerosis Self-Reported Case Prevalence Rates among</u>
<u>Females in GA, by County: Crude Rates as of November 2006</u>



<u>Figure 3:</u>
<u>Multiple Sclerosis Self-Reported Case Prevalence Rates among</u>
<u>Males in GA, by County: Crude Rates as of November 2006</u>




Table 2 shows the results of bivariate linear regression for self-reported prevalence and county level census data. We observed that the strength of the association between total MS rates and census data were largely driven by the female rates. Also, many of the variables show many highly significant values, especially per capita income. For these variables it was unlikely the association between the MS self-reported prevalence and the variable was due to chance alone, given the sample sizes.

Table	Table 2: Bivariate Linear Regression for Self-Reported-Case Prevalence and County Level Census Data.											
Variable	N	Total Rates	P-	R <sup>2</sup> of	Female	P-	R <sup>2</sup> of	Male rates	P-	$R^2$ of		
	(counties	and	value	the	Rates and	value	the	and	value	the		
	included in the	standardized	of beta <sup>a</sup>	total rate	standardized	of beta <sup>a</sup>	female Rate	standardized	of beta <sup>a</sup>	male rate		
	analysis)	beta (β)	(total	Tale	beta (β)	(female	Rate	beta (β)	male	Tale		
	ananyono,		rates)			rates)			rates)			
Total	159	0.259***	0.001	0.067	0.237**	0.003	0.056	0.175*	0.027	0.031		
population	100	0.200	0.001	0.007	0.207	0.000	0.000	0.170	0.027	0.001		
Percent female	159	0.143	0.073	0.02	0.098	0.217	0.010	0.142	0.075	0.020		
Female	159	0.260***	0.001	0.068	0.238**	0.003	0.056	0.176*	0.027	0.031		
population Male												
population	159	0.258**	0.001	0.067	0.237**	0.003	0.056	0.174*	0.028	0.03		
Percent white	159	0.036	0.655	0.001	0.007	0.934	0.000	0.085	0.284	0.007		
Percent												
black	159	-0.242**	0.002	0.058	-0.230**	0.003	0.053	-0.102	0.201	0.010		
Percent												
living in	159	-0.264***	0.001	0.070	-0.263***	0.001	0.069	-0.263***	0.001	0.069		
the same house												
Per capita	450	0.500***	0.000	0.070	0.450***	0.000	0.044	0.450***	0.000	0.044		
income	159	0.522****	0.000	0.272	0.459****	0.000	0.211	0.459****	0.000	0.211		
Square	159	-0.101	0.206	0.010	-0.116	0.147	0.013	-0.116	0.147	0.013		
miles land												

Table	2: Bivariate	e Linear Regres	ssion for	Self-Re	eported-Case P	revalence	and Cou	unty Level Cens	sus Data	ā.
Variable	N (counties included in the analysis)	Total Rates and standardized beta (β)	P- value of beta <sup>a</sup> (total rates)	R <sup>2</sup> of the total rate	Female Rates and standardized beta (β)	P- value of beta <sup>a</sup> (female rates)	R <sup>2</sup> of the female Rate	Male rates and standardized beta (β)	P- value of beta <sup>a</sup> male rates)	R <sup>2</sup> of the male rate
area										
People per square mile	159	0.262***	0.001	0.069	0.230**	0.004	0.053	0.230**	0.004	0.053

<sup>&</sup>lt;sup>a</sup> P-values are calculated using a two-tailed t distribution
\*Values are significant at the 0.05 level
\*\*Values are significant at the 0.01 level
\*\*\*Values are significant at the 0.001 level
\*\*\*\*Values are significant at the 0.0001 level

Table 3 shows the association between toxicants and environmental pollutants as county level emissions data and the total, female, and male self-reported-case prevalence rates of MS using bivariate linear regression. No statistically significant values were found for this table when using an alpha value of 0.05. However, two of the associations—Total rates and suspected cardiovascular or blood toxicants and suspected developmental toxicant—were very near to statistical significance.

Table 3: Biva	Table 3: Bivariate Linear Regression for Self-Reported-Case Prevalence versus County Level Emissions Data.											
Variable	N (counties included in the analysis)	Total rates and standard beta (β)	P-value of beta <sup>a</sup> (total rates)	R <sup>2</sup> of the total rate	Female rates and standard beta (β)	P-value of beta <sup>a</sup> (female rates)	R <sup>2</sup> of the female rate	Male rates and standard beta (β)	P-value of beta <sup>a</sup> male rates)	R <sup>2</sup> of the male rate		
Recognized carcinogens	83	-0.052	0.640	0.003	-0.064	0.563	0.004	0.002	0.987	0.000		
Suspected carcinogens	63	-0.087	0.496	800.0	-0.066	0.609	0.004	-0.029	0.822	0.001		
Suspected cardiovascular or blood toxicants	99	0.167	0.098	0.028	0.160	0.115	0.025	0.138	0.172	0.019		
Recognized development toxicants	79	0.196	0.084	0.038	0.181	0.111	0.033	0.112	0.325	0.013		
Suspected developmental toxicants	94	0.022	0.831	0.000	0.004	0.973	0.000	0.107	0.303	0.012		
Suspected endocrine toxicants	64	-0.049	0.705	0.002	0.001	0.991	0.000	-0.072	0.574	0.005		
Suspected immunotoxicants	96	0.056	0.586	0.003	0.087	0.401	0.007	0.016	0.877	0.000		

Table 3: Biva	riate Linear	Regressio	n for Self-	Reported-	-Case Prev	alence ve		nty Level Er	missions E	ata.
Variable	N	Total	P-value	R <sup>2</sup> of	Female	P-value	R <sup>2</sup> of	Male	P-value	R <sup>2</sup> of
	(counties	rates	of	the	rates	of beta <sup>a</sup>	the	rates	of	the
	included	and	beta <sup>a</sup>	total	and	(female	female	and	beta <sup>a</sup>	male
	in the	standard	(total	rate	standard	rates)	rate	standard	male	rate
	analysis)	beta (β)	rates)		beta (β)			beta (β)	rates)	
Suspected	100	0.073	0.472	0.005	0.058	0.564	0.003	0.097	0.338	0.009
kidney toxicants			-							
Suspected										
gastrointestinal	107	0.067	0.492	0.005	0.098	0.317	0.010	0.022	0.818	0.001
or liver toxicants										
Suspected										
muscoskeletal	32	-0.031	0.867	0.001	0.020	0.912	0.000	-0.081	0.658	0.007
toxicants										
Suspected	104	0.020	0.837	0.000	0.023	0.819	0.001	0.034	0.734	0.001
neurotoxicants										
Recognized		0.040				0.000		0.000	0.045	0.000
reproductive toxicants	70	-0.010	0.933	0.000	0.005	0.968	0.000	0.008	0.945	0.000
Suspected										
reproductive	103	0.030	0.767	0.001	0.036	0.721	0.001	0.036	0.721	0.001
toxicants	100	0.000	00.	0.00.	0.000	J.,	0.00	0.000	J.,	0.00.
Suspected										
respiratory	103	0.064	0.518	0.004	0.096	0.332	0.009	0.022	0.823	0.000
toxicants										
Suspected skin	404	0.055	0.500	0.000	0.000	0.070	0.000	0.040	0.000	0.000
or sense	101	0.055	0.586	0.003	0.090	0.372	0.008	0.010	0.920	0.000
toxicants										

Table 3: Biva	riate Linear	Regressio	n for Self-	Reported-	Case Prev	alence ve	rsus Coun	ty Level Er	nissions D	ata.
Variable	N	Total	P-value	R <sup>2</sup> of	Female	P-value	R <sup>2</sup> of	Male	P-value	R <sup>2</sup> of
	(counties	rates	of	the	rates	of beta <sup>a</sup>	the	rates	of	the
	included	and	beta <sup>a</sup>	total	and	(female	female	and	beta <sup>a</sup>	male
	in the	standard	(total	rate	standard	rates)	rate	standard	male	rate
	analysis)	beta (β)	rates)		beta (β)			beta (β)	rates)	
Acetaldehyde	14	-0.161	0.583	0.026	-0.202	0.489	0.041	0.061	0.835	0.004
Ammonia in	39	0.139	0.397	0.019	0.106	0.520	0.011	0.158	0.337	0.025
Formaldehyde	27	-0.138	0.493	0.019	-0.260	0.190	0.068	0.144	0.473	0.021
Lead/ lead	49	-0.146	0.318	0.021	-0.050	0.734	0.002	-0.164	0.260	0.027
compounds	10	0.110	0.010	0.021	0.000	0.701	0.002	0.101	0.200	0.027
Manganese	30	0.046	0.810	0.002	0.126	0.506	0.016	-0.062	0.744	0.004
compounds	00	0.010	0.010	0.002	0:120	0.000	0.010	0.002	0.7 1 1	0.001
Methanol	34	0.051	0.775	0.003	0.008	0.965	0.000	0.214	0.224	0.046
compounds		0.501	370	2.300	3.300	0.300	0.300	J.211	J. <b></b> .	3.310

The measurements for toxicants and selected produced compounds were reported in pounds for the year 2002.

<sup>&</sup>lt;sup>a</sup> P-values are calculated using a two-tailed t distribution

<sup>\*</sup>Values are significant at the 0.05 level

<sup>\*\*</sup>Values are significant at the 0.01 level

<sup>\*\*\*</sup>Values are significant at the 0.001 level

<sup>\*\*\*\*</sup>Values are significant at the 0.0001 level

Table 4 shows the associations between selected criteria air pollutants and overall MS self-reported-case prevalence using bivariate linear regression. The highest association and highest significance was observed for PM-10 and PM-2.5 mobile sources. Relatively weaker but still significant associations were observed with nitrogen oxides, carbon monoxide and volatile organic compounds. Finally, the association with these selected criteria air pollutants and female self-reported-case prevalence rates were generally stronger than those observed with male self-reported-case prevalence rates.

Table	Table 4: Bivariate Linear Regression: Total Self-Reported-Case Prevalence vs. Criteria Air Pollutant Data.										
Variable	N (counties	Total rates	P-value of beta <sup>a</sup>	R <sup>2</sup> of the	Female rates	P-value of beta <sup>a</sup>	R <sup>2</sup> of the	Male rates	P-value of beta <sup>a</sup>	R <sup>2</sup> of the	
	included in the analysis)	and standard beta (β)	(total rates)	total rate	and standard beta (β)	(female rates)	female rate	and standard beta (β)	male rates)	male rate	
Carbon Monoxide (mobile)	158	0.234**	0.003	0.055	0.208**	0.009	0.043	0.167*	0.036	0.028	
Carbon monoxide (area)	158	-0.078	0.330	0.006	-0.097	0.223	0.009	0.005	0.946	0.000	
Carbon monoxide (all)	158	0.193*	0.015	0.037	0.170*	0.033	0.029	0.144	0.072	0.021	
Nitrogen oxides (mobile)	158	0.242**	0.002	0.059	0.215**	0.007	0.046	0.176*	0.027	0.031	
Nitrogen oxides (area)	158	-0.003	0.974	0.000	0.03	0.705	0.001	-0.041	0.606	0.002	
Nitrogen oxides (all)	158	0.212**	0.007	0.045	0.213**	0.007	0.045	0.130	0.103	0.017	
PM-2.5 (mobile)	158	0.282***	0.000	0.079	0.264***	0.001	0.07	0.175*	0.028	0.030	

Table	Table 4: Bivariate Linear Regression: Total Self-Reported-Case Prevalence vs. Criteria Air Pollutant Data.										
Variable	N (counties included in the analysis)	Total rates and standard beta (β)	P-value of beta <sup>a</sup> (total rates)	R <sup>2</sup> of the total rate	Female rates and standard beta (β)	P-value of beta <sup>a</sup> (female rates)	R <sup>2</sup> of the female rate	Male rates and standard beta (β)	P-value of beta <sup>a</sup> male rates)	R <sup>2</sup> of the male rate	
PM-2.5 (area)	158	-0.093	0.247	0.009	-0.107	0.182	0.011	-0.001	0.992	0.000	
PM-2.5 (all)	158	0.107	0.179	0.012	0.073	0.363	0.005	0.142	0.076	0.020	
PM-10 (mobile)	158	0.276***	0.000	0.076	0.268***	0.001	0.072	0.159*	0.046	0.025	
PM-10 (area)	158	-0.045	0.578	0.002	-0.058	0.470	0.003	0.020	0.802	0.000	
PM-10 (all)	158	0.192*	0.016	0.037	0.179*	0.024	0.032	0.133	0.097	0.018	
Sulfur Dioxide (mobile)	158	0.136	0.089	0.018	0.118	0.141	0.014	0.136	0.089	0.018	
Sulfur Dioxide (area)	158	0.062	0.436	0.004	0.045	0.571	0.002	0.074	0.356	0.005	
Sulfur Dioxide (all)	158	0.083	0.297	0.007	0.105	0.189	0.011	0.034	0.675	0.001	
VOCs (mobile)	158	0.231**	0.003	0.054	0.206**	0.010	0.042	0.164*	0.039	0.027	
VOCs (area)	158	0.211**	0.008	0.045	0.196	0.013	0.039	0.138	0.083	0.019	

Table	Table 4: Bivariate Linear Regression: Total Self-Reported-Case Prevalence vs. Criteria Air Pollutant Data.										
Variable	N (counties included in the analysis)	Total rates and standard beta (β)	P-value of beta <sup>a</sup> (total rates)	R <sup>2</sup> of the total rate	Female rates and standard beta (β)	P-value of beta <sup>a</sup> (female rates)	R <sup>2</sup> of the female rate	Male rates and standard beta (β)	P-value of beta <sup>a</sup> male rates)	R <sup>2</sup> of the male rate	
VOCs (all)	158	0.243**	0.002	0.059	0.219**	0.006	0.048	0.174*	0.029	0.030	

<sup>&</sup>lt;sup>a</sup> p-values are calculated using a two-tailed t distribution

VOCs are volatile organic compounds. Also, the selected air pollutants were measured in 1999 and they were measured in tons of emissions to outdoor air.

<sup>\*</sup>Values are significant at the 0.05 level

<sup>\*\*</sup>Values are significant at the 0.01 level

<sup>\*\*\*</sup>Values are significant at the 0.001 level

<sup>\*\*\*\*</sup>Values are significant at the 0.0001 level

In table 5, using census, NEI and Total self-reported MS prevalence rates, the last model for stepwise multiple linear regression selected two characteristics, percapita income and PM-10 from area sources. Also, these two characteristics together had ability for this model to predict MS rates than per capita income alone. In these models, the beta coefficient was positive. This means as per capita income and PM-10 emissions to outdoor air from area sources increased, the total self-reported-case prevalence rates increased. Finally, the r-squared value estimated the ability for the model to predict the total self-reported-case prevalence rates by county. Therefore, in this model 26.2% of the total self-reported-case prevalence might have been explained by per capita income and PM-10 from area sources.

Table 5: Multivariate Linear Regression, Total MS Self-Repo	orted-Case Prevalence Rates a	nd Census and NEI	air pollutants
Model	Total rates and standardized beta (β)	P-value of beta <sup>a</sup> (total rates)	R <sup>2</sup> of the total rate
Per capita Income	0.512	0.000	0.262
Per Capita Income and PM-10 (area sources, 1999)	0.535	0.024	0.286
Per Capita Income	0.545	0.000	
PM-10 (area sources, 1999)	-0.158	0.024	

This analysis was done using stepwise multivariate linear regression with an acceptance p-value of 0.05 and an elimination p-value of 0.10. This analysis included only census data and selected criteria air pollutants.

As with the total rates, using census data and NEI, the final model summarized here for the female MS self-reported prevalence rates (table 6) selected per-capita income and PM-10 emissions to outdoor air from area sources for predicting female self-reported-case prevalence rates using multivariate linear regression. Also, the association was positive. The final model with the r-squared value of 0.225, suggesting 22.5% of the variance might be able to be explained using these two variables.

Table 6: Multivariate Linear Regression, Female MS Self-Reporte	ed-Case Prevalence Rates a	nd Census and NEI	air pollutants
	Total Rates and	P-value of Beta <sup>a</sup>	R <sup>2</sup> of the
Model	Standardized Beta (β)	(total rates)	Total Rate
Per capita Income	0.448	0.000	0.201
Per Capita Income and PM-10 (area sources, 1999)	0.474	0.030	0.225
Per Capita Income	0.481	0.000	
PM-10 (area sources, 1999)	-0.158	0.030	

This analysis was done using stepwise multivariate linear regression with an acceptance p-value of 0.05 and an elimination p-value of 0.10. This analysis included only census data and selected criteria air pollutants.

In table 7, using census data, NEI and male MS self-reported prevalence rates, stepwise multivariate linear regression selected only per capita income was included to estimate the self-reported-case prevalence of MS in males. The association calculated by the beta value was less than the association between per capita income and female or total rates. Lastly, the r-squared estimate was 0.114, or 11.4% of the variance in male MS rates might have been explained using this model.

Table 7: Multivariate Linear Regression, Male MS Self	-Reported-Case Prevalence Rates a	nd Census and NEI	air pollutants
	Total Rates and	P-value of Beta <sup>a</sup>	R <sup>2</sup> of the
Model	Standardized Beta (β)	(total rates)	Total Rate
Per capita Income	0.337	0.000	0.114
·			

This analysis was done using stepwise multivariate linear regression with an acceptance p-value of 0.05 and an elimination p-value of 0.10. This analysis included only census data and selected criteria air pollutants.

Table 8 shows steadily increasing odds ratios for total MS self-reported-case prevalence for several characteristics derived from U.S. Census data when separated by quartile: total population, male population, per capita income, and people per square mile. Also, the percent black and percent white show significantly strong but varying ORs. Also, people living in the same household in years 2000 and 2005 had ORs less than one.

Table 8: O	Table 8: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: Census Data and Total Self-Reported MS cases											
Variable	OR for lowest quartile (refer- ence)	OR for second quartile	95% CI <sup>a</sup> of the second quartile's OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile's OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the fourth quartile's OR	N (Number of people used for the OR)		
Total	4.00	4.04	(0.05.4.07)	004057	4.04	(0.07.4.00)	4500405	4 50****	(4.00.4.04)	7000070		
population	1.00	1.04	(0.85,1.27)	934957	1.04	(0.87,1.26)	1509185	1.52****	(1.28,1.81)	7208676		
Percent female	1.00	1.19****	(1.11,1.27)	5264143	1.04	(0.97,1.12)	4278364	0.93	(0.86,1.02)	3576459		
Female population	1.00	1.05	(0.86,1.29)	936425	1.09	(0.90,1.32)	1508117	1.56****	(1.31,1.85)	7209020		
Male population	1.00	1.02	(0.83,1.25)	936038	1.11	(0.92,1.34)	1510076	1.55****	(1.31,1.85)	7207260		
Percent white	1.00	0.85****	(0.78,0.92)	4517995	1.16****	(1.09,1.23)	6019018	0.89**	(0.82,0.96)	4929913		
Percent black	1.00	1.24***	(1.15,1.33)	4737748	0.89*	(0.81,0.98)	2890160	1.09*	(1.01,1.17)	4967140		

Table 8: O	dds Ratio	s(OR) and	95% Confide	ence Interva	al (CI) by (	Quartile: Cens	sus Data ai	nd Total S	elf-Reported	MS cases
Variable	OR for lowest quartile (refer- ence)	OR for second quartile	95% CI <sup>a</sup> of the second quartile's OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile's OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the fourth quartile's OR	N (Number of people used for the OR)
Percent living in the same house	1.00	0.69****	(0.64,0.74)	7847447	0.58****	(0.52,0.65)	6858190	0.62****	(0.54,0.70)	6697317
Per capita income	1.00	1.17	(0.97,1.41)	1293891	1.61****	(1.38,1.89)	2656723	2.22****	(1.92,2.58)	6158670
Square miles land	1.00	1.11**	(1.03,1.19)	4586763	1.01	(0.93,1.09)	3796684	0.94	(0.86,1.02)	3818753
People per sq. mile	1.00	1.27*	(1.05,1.56)	1099364	1.36***	(1.13,1.64)	1595362	1.90****	(1.60,2.26)	7085554

Each variable under analysis included 159 counties in GA.

<sup>&</sup>lt;sup>a</sup> 95% CI is calculated using the case-control study estimation model

<sup>\*</sup>Values are significant at the 0.05 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*</sup>Values are significant at the 0.01 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*\*</sup>Values are significant at the 0.001 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*\*\*</sup>Values are significant at the 0.0001 level using a Chi-squared test for 2 x 2 contingency tables

Table 9 shows total population, total female population, total male population, per capita income, and people per square mile had steadily increasing odds ratios for female MS self-reported-case prevalence across increasing quartiles. Also, as in Table 8, the percent living in the same house showed ORs less than one; however, this relationship did not continuously decrease across increasing quartiles for females.

Table 9:	Odds Rati	ios(OR) ar	nd 95% Confi	dence Inter	val (CI) by cases	/ Quartile: Ce	ensus Data	and Fema	le Self-Repo	rted MS
Variable	OR for lowest quartile (reference)	OR for second quartile	95% CI <sup>a</sup> of the second quartile OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the fourth quartile OR	N (Number of people used for the OR)
Total population	1.00	1.04	(0.82,1.33)	471185	1.05	(0.84,1.31)	758303	1.52***	(1.24,1.86)	3647621
Percent female	1.00	1.14*	(1.05,1.23)	2617130	0.99	(0.91,1.08)	2140378	0.89*	(0.81,0.99)	1801100
Female population	1.00	1.08	(0.85,1.38)	469590	1.11	(0.88,1.38)	759228	1.55****	(1.27,1.92)	3648235
Male population	1.00	1.03	(0.80,1.31)	473126	1.18	(0.94,1.48)	759731	1.58****	(1.29,1.95)	3645486
Percent white	1.00	0.88*	(0.80,0.97)	2307722	1.20****	(1.12,1.32)	3053523	0.95	(0.87,1.04)	2501961
Percent black	1.00	1.21****	(1.11,1.32)	2370527	0.87*	(0.77,0.97)	1450737	1.02	(0.94,1.12)	2522312
Percent living in same house	1.00	0.68****	(0.63,0.74)	3962140	0.61****	(0.54,0.70)	3457814	0.59****	(0.51,0.69)	3376835

Table 9:	Table 9: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: Census Data and Female Self-Reported MS cases									
Variable	OR for lowest quartile (reference)	OR for second quartile	95% CI <sup>a</sup> of the second quartile OR	N (Number of people used for the OR)	OR for third quartile	95% Cl <sup>a</sup> of the third quartile OR	N (Number of people used for the OR)	OR for highest quartile	95% Cl <sup>a</sup> of the fourth quartile OR	N (Number of people used for the OR)
Per capita income	1.00	1.21	(0.97,1.51)	648920	1.65****	(1.37,2.00)	1346989	2.23****	(1.86,2.67)	3098085
Square miles land	1.00	1.12*	(1.03,1.22)	2331699	1.07	(0.98,1.18)	1911463	0.96	(0.87,1.06)	1934333
People per sq. mile	1.00	1.22	(0.96,1.54)	554040	1.38**	(1.11,1.71)	802485	1.85***	(1.51,2.27)	3585499

Each variable under analysis included 159 counties in GA.

<sup>&</sup>lt;sup>a</sup> 95% CI is calculated using the case-control study estimation model

<sup>\*</sup>Values are significant at the 0.05 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*</sup>Values are significant at the 0.01 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*\*</sup>Values are significant at the 0.001 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*\*\*</sup>Values are significant at the 0.0001 level using a Chi-squared test for 2 x 2 contingency tables

Table 10 shows MS self-reported-case prevalence for the male population showed no consistent results; however, the total population, the female population, the male population, and the per capita income showed increasing odds ratios starting from the second quartile. Also, the percent living in the same house seems to have a protective effect, i.e. OR and CI consistently less than 1.00.

Table 10: 0	Odds Rati	os(OR) ar	nd 95% Cor	nfidence Int	` ,	by Quartile:	Census Da	ta and Ma	ale Self-Rep	orted MS
					cases			1		
Variable	OR for	OR for	95% Cl <sup>a</sup>	N	OR for	95% Cl <sup>a</sup>	N	OR for	95% Cl <sup>a</sup>	N
	lowest	second	of the	(Number	third	of the	(Number	highest	of the	(Number
	quartile	quartile	second	of	quartile	third	of	quartile	fourth	of
	(refer-		quartile's	people		quartile's	people		quartile's	people
	ence)		OR	used for		OR	used for		OR	used for
				the OR)			the OR)			the OR)
Total			(0.58,			(0.65,			(1.01,	
population	1.00	0.89	1.37)	463772	0.97	1.43)	750882	1.43*	2.05)	3561055
Percent			(1.06,			(0.93,			(0.82,	
female	1.00	1.22**	1.41)	2647012	1.1	1.29)	2137986	0.99	1.18)	1775359
			•			•			•	
Female			(0.55,			(0.66,			(1.01,	
population	1.00	0.85	1.31)	466835	0.98	1.45)	748889	1.44*	2.06)	3560785
Male			(0.56,			(0.58,			(0.96,	
population	1.00	0.85	1.30)	462912	0.86	1.15)	750345	1.36	1.92)	3561774
Percent			(0.63,			(0.93,			(0.65,	
white	1.00	0.75**	0.90)	2210272	1.06	1.20)	2965495	0.76***	0.89)	2427952
Percent			(1.10,			(0.81,			(1.08,	
black	1.00	1.29**	1.51)	2367221	1.01	1.24)	1439423	1.27**	1.49)	2444828
Percent										
living in										
same			(0.60,			(0.38,			(0.53,	
house	1.00	0.70****	0.82)	3885306	0.50****	0.65)	3400375	0.69***	0.90)	3320481

Table 10:	Odds Rati	os(OR) ar	nd 95% Cor	nfidence Int	erval (CI)	by Quartile:	Census Da	ta and Ma	ile Self-Rep	orted MS
					cases					
Variable	OR for lowest quartile (reference)	OR for second quartile	95% CI <sup>a</sup> of the second quartile's OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile's OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the fourth quartile's OR	N (Number of people used for the OR)
Per capita income	1.00	0.96	(0.65, 1.42)	644971	1.41*	(1.03, 1.95)	1309734	2.04***	(1.51, 2.77)	3060585
Square miles land area 2000	1.00	1.13	(0.97, 1.32)	2255064	0.90	(0.76, 1.07)	1885221	0.92	(0.78, 1.09)	1884420
People per sq. mile	1.00	1.43	(0.93, 2.20)	545324	1.26	(0.84, 1.91)	792877	2.01***	(1.38, 2.94)	3500055

Each variable under analysis includes 159 counties in GA.

<sup>&</sup>lt;sup>a</sup> 95% CI is calculated using the case-control study estimation model

<sup>\*</sup>Values are significant at the 0.05 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*</sup>Values are significant at the 0.01 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*\*</sup>Values are significant at the 0.001 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*\*\*</sup>Values are significant at the 0.0001 level using a Chi-squared test for 2 x 2 contingency tables

Table 11 showed increases in the odds ratios for total MS self-reported cases, and tended to increase from the second to the fourth quartiles for county level emissions to outdoor air recognized developmental toxicants, suspected endocrine toxicants, suspected immunotoxicants, suspected kidney toxicants and ammonia. For many of these variables, only the fourth quartile showed statistical significance. Also, there seemed to be statistically significant lower odds ratios for recognized carcinogens.

Table 11: Od	dds Ratios	(OR) and	95% Confid		` '		County Leve	I Emission	ns and Tota	al Self-
	1	T			d MS case					
Variable (# of counties in analysis for each variable <sup>b</sup> )	OR for lowest quartile (reference)	OR for second quartile	95% Cl <sup>a</sup> of the second quartile OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the highest quartile OR	N (Number of people used for the OR)
Recognized Carcinogens (83)	1.00	0.43****	(0.40, 0.46)	2683293	0.46****	(0.43, 0.49)	3977208	0.44****	(0.41, 0.48)	3188848
Suspected Carcinogens (63)	1.00	1.15**	(1.05, 1.27)	2800765	0.88*	(0.79, 0.98)	2112213	1.15**	(1.05, 1.25)	4245365
Suspected Cardio- vascular or Blood Toxicants (99)	1.00	1.09	(0.97, 1.23)	1796302	1.04	(0.93, 1.15)	2975051	1.19***	(1.08, 1.31)	4844048

Table 11: O	dds Ratios	(OR) and	95% Confid		, , ,		County Leve	I Emission	ns and Tot	al Self-
Variable (# of counties in analysis for each variable <sup>b</sup> )	OR for lowest quartile (reference)	OR for second quartile	95% Cl <sup>a</sup> of the second quartile OR	N (Number of people used for the OR)	d MS case OR for third quartile	95% Cl <sup>a</sup> of the third quartile OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the highest quartile OR	N (Number of people used for the OR)
Recognized Develop- mental toxicants (79)	1.00	0.90*	(0.81, 1.00)	2358067	1.04	(0.95, 1.15)	2989780	1.08	(0.98, 1.18)	3881857
Suspected Develop- mental toxicants (94)	1.00	1.04	(0.94, 1.15)	2249010	0.92	(0.84, 1.02)	3073401	1.16***	(1.06, 1.27)	4435056
Suspected Endocrine toxicants (64)	1.00	0.92	(0.83, 1.02)	2208012	0.95	(0.87, 1.04)	3074432	1.15***	(1.06, 1.25)	4151693
Suspected Immuno- toxicants (96)	1.00	0.74***	(0.67, 0.82)	2319209	0.96	(0.87, 1.04)	3165315	1.10*	(1.02, 1.20)	4769708

Table 11: Od	dds Ratios	(OR) and	95% Confid				County Leve	I Emission	ns and Tota	al Self-
				Reported	d MS case	es es				
Variable (# of counties in analysis for each variable <sup>b</sup> )	OR for lowest quartile (reference)	OR for second quartile	95% Cl <sup>a</sup> of the second quartile OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the highest quartile OR	N (Number of people used for the OR)
Suspected Kidney toxicants (100)	1.00	1.03	(0.91, 1.15)	1986463	1.07	(0.96, 1.19)	3024089	1.26****	(1.15, 1.39)	4653514
Suspected Gastro- intestinal or liver toxicants (107)	1.00	1.20**	(1.08, 1.35)	2096331	1.04	(0.93, 1.15)	3101108	1.34****	(1.21, 1.47)	4699751
Suspected Musco- skeletal toxicants (32)	1.00	1.09	(0.99, 1.19)	2601356	0.96	(0.87, 1.07)	2387914	1.29****	(1.18, 1.41)	2617585
Suspected Neuro- toxicants (104)	1.00	1.05	(0.95, 1.15)	2766274	0.86**	(0.78, 0.95)	2611959	1.16***	(1.06, 1.26)	4723348

Table 11: Od	dds Ratios	(OR) and	95% Confid		al (CI) by d d MS case		ounty Leve	l Emission	s and Tot	al Self-
Variable (# of counties in analysis for each variable <sup>b</sup> )	OR for lowest quartile (reference)	OR for second quartile	95% CI <sup>a</sup> of the second quartile OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the highest quartile OR	N (Number of people used for the OR)
Recognized Reproductive toxicants (70)	1.00	1.03	(0.93, 1.14)	2506688	0.86*	(0.77, 0.97)	1965624	1.16***	(1.06, 1.27)	4299971
Suspected Reproductive toxicants (103)	1.00	1.20***	(1.08, 1.34)	2265535	1.08	(0.97, 1.19)	2981303	1.32****	(1.20, 1.45)	4734854
Suspected Respiratory Toxicants (103)	1.00	1.04	(0.94, 1.16)	2109269	0.92	(0.84, 1.01)	3229846	1.21****	(1.11, 1.31)	4797755
Suspected Skin or Sense Toxicants (101)	1.00	1.03	(0.93, 1.14)	2222602	0.90*	(0.81, 0.99)	3063788	1.19****	(1.09, 1.29)	4764846

Table 11: Od	Table 11: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: County Level Emissions and Total Self- Reported MS cases									
				Reported	d MS case	es				
Variable (# of counties in analysis for	OR for lowest quartile	OR for second	95% CI <sup>a</sup> of the second	N (Number of people	OR for third quartile	95% CI <sup>a</sup> of the	N (Number of	OR for highest quartile	95% Cl <sup>a</sup> of the	N (Number of
each variable <sup>b</sup> )	(refer- ence)	quartile	quartile OR	used for the OR)	quartile	third quartile OR	people used for the OR)	quartile	highest quartile OR	people used for the OR)
Acetaldehyde (14)	1.00	0.75	(0.55, 1.03)	277007	0.88	(0.72, 1.09)	692052	0.91	(0.73, 1.14)	513424
Ammonia (39)	1.00	0.91	(0.78, 1.05)	1684506	1.16**	(1.05, 1.27)	2564465	1.15**	(1.06, 1.25)	3195643
Formaldehyde (27)	1.00	1.45**	(1.14, 1.84)	472234	1.32**	(1.08, 1.61)	999745	1.53****	(1.26, 1.84)	1410500
Lead/ lead compounds (49)	1.00	0.92	(0.81, 1.03)	1777914	0.83**	(0.74, 0.93)	1947538	1.14**	(1.04, 1.25)	2454203
Manganese compounds (30)	1.00	1	(0.87, 1.16)	1819989	1.32***	(1.14, 1.53)	1313616	0.96	(0.82, 1.14)	1009772
Methanol (34)	1.00	1.14	(0.99, 1.31)	1642066	1.12	(0.97, 1.30)	1334797	1.05	(0.91, 1.21)	1336137

Table 11: Od	dds Ratios	(OR) and	95% Confid		` '		County Leve	I Emission	ns and Tot	al Self-				
	Reported MS cases													
Variable (# of counties in analysis for each variable <sup>b</sup> )	OR for lowest quartile (reference)	OR for second quartile	95% Cl <sup>a</sup> of the second quartile OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the highest quartile OR	N (Number of people used for the OR)				

The measurements for toxicants and selected produced compounds were reported in pounds for the year 2002.

<sup>&</sup>lt;sup>a</sup> 95% CI is calculated using the case-control study estimation model.

<sup>&</sup>lt;sup>b</sup> Every county that had information concerning the respective variable was included. The total number of counties in GA is 159 at the time of this study.

<sup>\*</sup>Values are significant at the 0.05 level using a Chi-squared test for 2 x 2 contingency tables.

<sup>\*\*</sup>Values are significant at the 0.01 level using a Chi-squared test for 2 x 2 contingency tables.

<sup>\*\*\*</sup>Values are significant at the 0.001 level using a Chi-squared test for 2 x 2 contingency tables.

<sup>\*\*\*\*</sup>Values are significant at the 0.0001 level using a Chi-squared test for 2 x 2 contingency tables.

Table 12 showed small to modest increases in the odds ratios for female MS self-reported cases, and tended to increase from the second to the fourth quartiles of county level emissions in pounds to outdoor air recognized developmental toxicants, suspected endocrine toxicants, suspected immunotoxicants, suspected kidney toxicants and ammonia. Also, there seemed to be lower odds ratios for recognized carcinogens; the values were similar across quartiles.

Table 12: Odd	ds Ratios(C	OR) and 95	% Confide		l (CI) by Qu d MS cases		unty Level E	Emissions	and Fema	ale Self-
Variable (# of counties in analysis <sup>b</sup> )	OR for lowest quartile (reference)	OR for second quartile	95% CI <sup>a</sup> of the second quartile' s OR	N (Number of people used for the OR)	OR for third quartile	95% Cl <sup>a</sup> of the third quartile' s OR	N (Number of people used for the OR)	OR for highest quartile	95% Cl <sup>a</sup> of the highest quartil e's OR	N (Number of people used for the OR)
Recognized Carcinogens (83)	1.00	0.86**	(0.77, 0.96)	1345737	0.89*	(0.81, 0.98)	2004536	0.85**	(0.77, 0.94)	1626595
Suspected Carcinogens (63)	1.00	1.14*	(1.02, 1.27)	1428998	0.92	(0.81, 1.05)	1080212	1.12*	(1.01, 1.24)	2155182
Suspected Cardio- vascular or Blood Toxicants (99)	1.00	1.02	(0.92, 1.21)	908939	1.04	(0.92, 1.18)	1499244	1.16**	(1.04, 1.31)	2457076
Recognized Develop- mental toxicants (79)	1.00	0.92	(0.81, 1.04)	1181017	1.01	(0.91, 1.14)	1513394	1.04	(0.93, 1.16)	1979912
Suspected Develop- mental toxicants (94)	1.00	1.04	(0.92, 1.17)	1140228	0.94	(0.84, 1.06)	1544766	1.12	(1.01, 1.24)	2255240

Table 12: Odd	ds Ratios(C	R) and 95	% Confide		(CI) by Qu		ınty Level E	missions	and Fema	ale Self-
Variable (# of counties in analysis <sup>b</sup> )	OR for lowest quartile (reference)	OR for second quartile	95% CI <sup>a</sup> of the second quartile' s OR	N (Number of people used for the OR)	OR for third quartile	95% Cl <sup>a</sup> of the third quartile' s OR	N (Number of people used for the OR)	OR for highest quartile	95% Cl <sup>a</sup> of the highest quartil e's OR	N (Number of people used for the OR)
Suspected Endocrine toxicants (64)	1.00	0.94	(0.83, 1.07)	1126196	1.01	(0.90, 1.12)	1551054	1.14**	(1.04, 1.26)	2109476
Suspected Immuno- toxicants (96)	1.00	0.72***	(0.64, 0.82)	1175080	0.96	(0.86, 1.06)	1584123	1.07	(0.97, 1.17)	2423900
Suspected Kidney toxicants (100)	1.00	1.04	(0.90, 1.19)	1006583	1.10	(0.97, 1.24)	1516143	1.24***	(1.10, 1.39)	2366472
Suspected Gastro- intestinal or liver toxicants (107)	1.00	1.15*	(1.01, 1.31)	1056961	1.05	(0.93, 1.19)	1563048	1.28****	(1.15, 1.44)	2390813
Suspected Musco-skeletal toxicants (32)	1.00	1.03	(0.92, 1.15)	1311624	0.96	(0.85, 1.09)	1214744	1.29****	(1.16, 1.43)	1313325
Suspected Neuro- toxicants (104)	1.00	1.07	(0.95, 1.20)	1385840	0.89	(0.79, 1.01)	1323064	1.14*	(1.03, 1.26)	2398195

Table 12: Odd	Table 12: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: County Level Emissions and Female Self-Reported MS cases									
Variable (# of counties in analysis <sup>b</sup> )	OR for lowest quartile (reference)	OR for second quartile	95% Cl <sup>a</sup> of the second quartile' s OR	N (Number of people used for the OR)	OR for third quartile	95% Cl <sup>a</sup> of the third quartile' s OR	N (Number of people used for the OR)	OR for highest quartile	95% Cl <sup>a</sup> of the highest quartil e's OR	N (Number of people used for the OR)
Recognized Reproductive toxicants (70)	1.00	1.05	(0.93, 1.18)	1257874	0.88	(0.77, 1.01)	993398	1.13*	(1.01, 1.25)	2182681
Suspected Reproductive toxicants (103)	1.00	1.15*	(1.01, 1.31)	1144559	1.09	(0.97, 1.23)	1498011	1.25****	(1.12, 1.40)	2408274
Suspected Respiratory Toxicants (103)	1.00	1.05	(0.93, 1.19)	1065348	0.95	(0.85, 1.06)	1624733	1.19***	(1.07, 1.32)	2438482
Suspected Skin or Sense Toxicants (101)	1.00	1.03	(0.91, 1.16)	1124204	0.94	(0.84, 1.05)	1538373	1.18*	(1.06, 1.30)	2421997
Acetaldehyde (14)	1.00	0.9	(0.62, 1.30)	138622	0.89	(0.69, 1.15)	358164	1.08	(0.83, 1.40)	262337
Ammonia (39)	1.00	0.91	(0.76, 1.08)	848900	1.13*	(1.01, 1.26)	1289724	1.11*	(1.00, 1.23)	1621452

Table 12: Odd	ds Ratios(C	OR) and 95	5% Confide		(CI) by Qu		ınty Level E	missions	and Fema	ale Self-
Variable (# of counties in analysis <sup>b</sup> )	OR for lowest quartile (reference)	OR for second quartile	95% CI <sup>a</sup> of the second quartile' s OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile' s OR	N (Number of people used for the OR)	OR for highest quartile	95% Cl <sup>a</sup> of the highest quartil e's OR	N (Number of people used for the OR)
Formaldehyde (27)	1.00	1.42**	(1.08, 1.87)	238691	1.23	(0.98, 1.55)	512801	1.34**	(1.08, 1.67)	720232
Lead/ lead compounds (49)	1.00	0.93	(0.81, 1.07)	891840	0.81*	(0.71, 0.93)	985396	1.15*	(1.03, 1.28)	1226351
Manganese compounds (30)	1.00	1.03	(0.87, 1.23)	919813	1.34***	(1.12, 1.60)	662482	1.02	(0.84, 1.24)	515580
Methanol (34)	1.00	1.13	(0.96, 1.34)	837831	1.08	(0.91, 1.28)	681676	1.07	(0.90, 1.28)	689674

The measurements for toxicants and selected produced compounds were reported in pounds for the year 2002.

<sup>&</sup>lt;sup>a</sup> 95% CI is calculated using the case-control study estimation model.

<sup>&</sup>lt;sup>b</sup> Every county that had information concerning the respective variable was included. The total number of counties in GA is 159 at the time of this study.

<sup>\*</sup>Values are significant at the 0.05 level using a Chi-squared test for 2 x 2 contingency tables.

<sup>\*\*</sup>Values are significant at the 0.01 level using a Chi-squared test for 2 x 2 contingency tables.

<sup>\*\*\*</sup>Values are significant at the 0.001 level using a Chi-squared test for 2 x 2 contingency tables.

<sup>\*\*\*\*</sup>Values are significant at the 0.0001 level using a Chi-squared test for 2 x 2 contingency tables.

Table 13 showed odds ratios tended to increase for male MS self-reported from the second to the fourth quartiles of county level emissions in pounds of outdoor air for recognized developmental toxicants, suspected endocrine toxicants, suspected immunotoxicants, suspected kidney toxicants, ammonia and formaldehyde. It is important to note only the counties that reported values for the variable of interest were included for each variable.

Table 13: O	dds Ratios	(OR) and 9	95% Confid	lence Interva	al (CI) by ( MS case		ounty Level	Emission	s and Male	e Self-
Variable (# of counties in analysis <sup>b</sup> )	OR for lowest quartile	OR for second quartile	95% CI <sup>a</sup> of the second quartile OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the highest quartile OR	N (Number of people used for the OR)
Recognized Carcinogens (83)	1.00	0.84	(0.69, 1.03)	1336736	0.97	(0.81, 1.15)	1971852	0.95	(0.79, 1.15)	1561433
Suspected Carcinogens (63)	1.00	1.19	(0.97, 1.47)	1371767	0.88	(0.69, 1.13)	1032001	1.31**	(1.08, 1.58)	2090183
Suspected Cardio- vascular or Blood Toxicants (99)	1.00	1.25	(0.97, 1.61)	887363	1.08	(0.85, 1.36)	1475807	1.32*	(1.06, 1.64)	2386972
Recognized Develop- mental toxicants (79)	1.00	0.95	(0.75, 1.20)	1177051	1.20	(0.97, 1.49)	1476386	1.25*	(1.02, 1.53)	1901945
Suspected Develop- mental toxicants (94)	1.00	1.05	(0.83, 1.32)	1108782	0.90	(0.72, 1.11)	1528635	1.31**	(1.09, 1.59)	2179816

Table 13: O	Table 13: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: County Level Emissions and Male Self-Reported MS cases									
Variable (# of counties in analysis <sup>b</sup> )	OR for lowest quartile	OR for second quartile	95% Cl <sup>a</sup> of the second quartile OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the highest quartile OR	N (Number of people used for the OR)
Suspected Endocrine toxicants (64)	1.00	0.90	(0.72, 1.13)	1081817	0.93	(0.76, 1.12)	1523378	1.22*	(1.02, 1.46)	2042218
Suspected Immuno- toxicants (96)	1.00	0.88	(0.70, 1.11)	1144129	0.99	(0.81, 1.21)	1581192	1.27**	(1.06, 1.52)	2345808
Suspected Kidney toxicants (100)	1.00	1.02	(0.79, 1.32)	979880	1.06	(0.84, 1.33)	1507946	1.33**	(1.07, 1.65)	2287042
Suspected Gastro- intestinal or liver toxicants (107)	1.00	1.31*	(1.03, 1.66)	1039370	1.01	(0.80, 1.27)	1538060	1.42***	(1.15, 1.75)	2308938
Suspected Muscoskeletal toxicants (32)	1.00	1.21*	(1.00, 1.47)	1288785	0.90	(0.72, 1.13)	1172223	1.22*	(1.01, 1.47)	1303313
Suspected Neuro- toxicants (104)	1.00	0.97	(0.80, 1.19)	1380434	0.72**	(0.58, 0.90)	1288895	1.15	(0.96, 1.38)	2325153

Table 13: O	Table 13: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: County Level Emissions and Male Self-Reported MS cases									
Variable (# of counties in analysis <sup>b</sup> )	OR for lowest quartile	OR for second quartile	95% Cl <sup>a</sup> of the second quartile OR	N (Number of people used for the OR)	OR for third quartile	95% Cl <sup>a</sup> of the third quartile OR	N (Number of people used for the OR)	OR for highest quartile	95% Cl <sup>a</sup> of the highest quartile OR	N (Number of people used for the OR)
Recognized Reproductive toxicants (70)	1.00	1.13	(0.90, 1.42)	1248814	0.96	(0.74, 1.24)	972226	1.44***	(1.18, 1.76)	2117290
Suspected Reproductive toxicants (103)	1.00	1.40**	(1.10, 1.79)	1120976	1.08	(0.85, 1.37)	1483292	1.51***	(1.22, 1.87)	2326580
Suspected Respiratory Toxicants (103)	1.00	0.98	(0.78, 1.23)	1043921	0.86	(0.70, 1.06)	1605113	1.21*	(1.01, 1.45)	2359273
Suspected Skin or Sense Toxicants (101)	1.00	0.97	(0.78, 1.21)	1098398	0.80*	(0.65, 0.98)	1525415	1.16	(0.97, 1.39)	2342849
Acetaldehyde (14)	1.00	0.62	(0.29, 1.30)	138385	1.03	(0.66, 1.60)	333888	0.77	(0.47, 1.27)	251087
Ammonia (39)	1.00	0.82	(0.58, 1.15)	835606	1.17	(0.96, 1.43)	1274741	1.16	(0.97, 1.39)	1574191

Table 13: O	dds Ratios	(OR) and	95% Confid	lence Interva	al (CI) by 0 MS case		ounty Level	Emission	s and Mal	e Self-
Variable (# of counties in analysis <sup>b</sup> )	OR for lowest quartile	OR for second quartile	95% Cl <sup>a</sup> of the second quartile OR	N (Number of people used for the OR)	OR for third quartile	95% Cl <sup>a</sup> of the third quartile OR	N (Number of people used for the OR)	OR for highest quartile	95% Cl <sup>a</sup> of the highest quartile OR	N (Number of people used for the OR)
Formaldehyde (27)	1.00	1.90*	(1.06, 3.43)	233543	1.86**	(1.13, 3.10)	486944	2.38***	(1.48, 3.87)	690268
Lead/ lead compounds (49)	1.00	0.88	(0.67, 1.16)	886074	0.93	(0.72, 1.19)	962142	1.17	(0.96, 1.42)	1227852
Manganese compounds (30)	1.00	1.06	(0.77, 1.47)	900176	1.37	(0.99, 1.91)	651134	1.03	(0.72, 1.48)	494192
Methanol (34)	1.00	1.12	(0.83, 1.51)	804235	1.20	(0.88, 1.63)	653121	1.04	(0.76, 1.42)	646463

The measurements for toxicants and selected produced compounds were reported in pounds for the year 2002.

<sup>&</sup>lt;sup>a</sup> 95% CI is calculated using the case-control study estimation model.

<sup>&</sup>lt;sup>b</sup> Every county that had information concerning the respective variable was included. The total number of counties in GA is 159 at the time of this study.

<sup>\*</sup>Values are significant at the 0.05 level using a Chi-squared test for 2 x 2 contingency tables.

<sup>\*\*</sup>Values are significant at the 0.01 level using a Chi-squared test for 2 x 2 contingency tables.

<sup>\*\*\*</sup>Values are significant at the 0.001 level using a Chi-squared test for 2 x 2 contingency tables.

<sup>\*\*\*\*</sup>Values are significant at the 0.0001 level using a Chi-squared test for 2 x 2 contingency tables.

Table 14 shows increasing odds ratios for total MS self-reported cases with increasing emissions of selected criteria air pollutants to outdoor air for carbon monoxide (mobile sources), nitrogen oxides (mobile and areas sources), PM-2.5 (mobile sources), PM-10 (mobile sources), and volatile organic compounds (mobile sources). In addition, the odds ratios tended to increase from the second to the fourth quartile for nitrogen (all sources), PM-2.5 (area and all sources), PM-10 (area and all sources), sulfur dioxide (area and all sources), and volatile organic compounds (area sources). The majority of these findings were highly statistically significant.

Table 1	14: Odds	Ratios(OR	) and 95%		Interval (	, •	rtile: Criteria	a Air Pollut	tants and T	otal Self-
Variable	OR for lowest quartile	OR for second quartile	95% CI <sup>a</sup> of the second quartile OR	N (Number of people used for the OR)	OR for third quartile	95% Cl <sup>a</sup> of the third quartile OR	N (Number of people used for the OR)	OR for highest quartile	95% Cl <sup>a</sup> of the fourth quartile OR	N (Number of people used for the OR)
Carbon Monoxide (mobile)	1.00	1.21*	(1.00, 1.47)	1067078	1.26*	(1.11, 1.42)	1546590	1.63****	(1.38, 1.92)	6946275
Carbon monoxide (area)	1.00	1.29****	(1.15, 1.44)	2906874	1.26****	(1.12, 1.42)	2366165	1.46****	(1.31, 1.62)	5166698
Carbon monoxide (all)	1.00	0.75****	(0.65, 0.86)	1343172	0.72****	(0.64, 0.82)	1915213	1.04	(0.94, 1.15)	6850832
Nitrogen oxides (mobile)	1.00	1.01	(0.85, 1.19)	1205793	1.04	(0.89, 1.21)	1596924	1.40****	(1.22, 1.61)	6930996
Nitrogen oxides (area)	1.00	1.02	(0.90, 1.15)	1849188	1.09	(0.97, 1.22)	2364973	1.23****	(1.11, 1.36)	6178314
Nitrogen oxides (all)	1.00	0.94	(0.79, 1.12)	1128444	0.94	(0.80, 1.11)	1702089	1.43****	(1.23, 1.65)	6793742
PM-2.5 (mobile)	1.00	1.21	(0.99, 1.49)	993506	1.22*	(1.00, 1.48)	1510700	1.75****	(1.47, 2.10)	6998801
PM-2.5	1.00	0.94	(0.85,	2858242	1.01	(0.90,	2444023	1.16**	(1.05,	5065518

Table '	14: Odds	Ratios(OR	R) and 95%	Confidence Re	Interval ( ported MS	. •	rtile: Criteria	a Air Pollu	tants and T	otal Self-
Variable	OR for lowest quartile	OR for second quartile	95% Cl <sup>a</sup> of the second quartile OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the fourth quartile OR	N (Number of people used for the OR)
(area)			1.05)			1.12)			1.28)	
PM-2.5 (all)	1.00	0.89	(0.77, 1.03)	1384414	1.00	(0.88, 1.13)	2425057	1.23***	(1.09, 1.38)	6163914
PM-10 (mobile)	1.00	1.00	(0.84, 1.20)	1048779	1.06	(0.90, 1.25)	1740213	1.47****	(1.27, 1.70)	6858117
PM-10 (area)	1.00	0.93	(0.83, 1.03)	2475108	0.93	(0.84, 1.04)	2493041	1.12*	(1.02, 1.24)	5468954
PM-10 (all)	1.00	0.87	(0.74, 1.01)	1172668	0.87*	(0.76, 0.99)	2312006	1.20**	(1.06, 1.35)	6359593
Sulfur dioxide (mobile)	1.00	1.09	(0.91, 1.30)	1077110	0.95	(0.81, 1.13)	1598051	1.50****	(1.29, 1.74)	6960012
Sulfur dioxide (area)	1.00	0.93	(0.79, 1.09)	1256009	0.94	(0.81, 1.09)	1728222	1.43****	(1.27, 1.62)	6942742
Sulfur dioxide (all)	1.00	0.94	(0.79, 1.12)	1191739	1.15	(0.98, 1.34)	2034489	1.36****	(1.18, 1.57)	6396515

Table '	14: Odds	Ratios(OF	R) and 95%		Interval ( ported MS	. •	rtile: Criteria	a Air Pollu	tants and T	otal Self-
Variable	OR for lowest quartile	OR for second quartile	95% CI <sup>a</sup> of the second quartile OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the fourth quartile OR	N (Number of people used for the OR)
VOCs (mobile)	1.00	1.09	(0.91, 1.30)	1087960	1.14	(0.96, 1.34)	1593661	1.48***	(1.27, 1.72)	6939764
VOCs (area)	1.00	0.80*	(0.68, 0.95)	1092424	0.94	(0.81, 1.09)	1727586	1.25***	(1.09, 1.42)	6888893
VOCs (all)	1.00	1.10	(0.92, 1.33)	1087682	1.09	(0.92, 1.30)	1546984	1.54***	(1.32, 1.80)	6962477

Each variable under analysis included 158 of the 159 counties in GA. VOCs are volatile organic compounds, and PM is particulate matter. Also, the selected air pollutants were reported as emissions in tons, in 1999.

<sup>&</sup>lt;sup>a</sup> 95% CI is calculated using the case-control study estimation model

<sup>\*</sup>Values are significant at the 0.05 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*</sup>Values are significant at the 0.01 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*\*</sup>Values are significant at the 0.001 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*\*\*</sup>Values are significant at the 0.0001 level using a Chi-squared test for 2 x 2 contingency tables

Table 15 shows increasing odds ratios for female MS self-reported cases with increasing emissions of selected criteria air pollutants to outdoor air for carbon monoxide (mobile and area sources), nitrogen oxides (areas sources), and PM-2.5 (mobile sources). In addition, the odds ratios tended to increase as we go from the second to the fourth quartile for nitrogen oxides (mobile and all sources), PM-2.5 (area and all sources), PM-10 (mobile, area and all sources), sulfur dioxide (area and all sources), and volatile organic compounds (mobile and area sources). The majority of the findings were highly statistically significant.

Table 15	Table 15: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: Criteria Air Pollutants and Female Self-Reported MS cases									
Variable	OR for lowest quartile	OR for second quartile	95% Cl <sup>a</sup> of the second quartile OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile's OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the fourth quartile's OR	N (Number of people used for the OR)
Carbon Monoxide (mobile)	1.00	1.13	(0.90, 1.41)	535442	1.23	(0.99, 1.51)	776385	1.55****	(1.28, 1.88)	3515655
Carbon monoxide (area)	1.00	1.26***	(1.10, 1.44)	1478302	1.27***	(1.10, 1.46)	1192659	1.44***	(1.27, 1.63)	2604139
Carbon monoxide (all)	1.00	0.76***	(0.64, 0.89)	676178	0.72****	(0.62, 0.83)	965122	1.04	(0.92, 1.17)	3469210
Nitrogen oxides (mobile)	1.00	0.98	(0.81, 1.19)	604846	1.01	(0.84, 1.21)	803584	1.34***	(1.15, 1.58)	3508701
Nitrogen oxides (area)	1.00	1.02	(0.88, 1.18)	931641	1.08	(0.95, 1.24)	1198135	1.23***	(1.10, 1.39)	3128221
Nitrogen oxides (all)	1.00	0.87	(0.71, 1.07)	565409	0.90	(0.75, 1.09)	858482	1.35***	(1.14, 1.60)	3439471
PM-2.5 (mobile)	1.00	1.13	(0.88, 1.44)	498241	1.19	(0.95, 1.49)	760421	1.68****	(1.37, 2.07)	3543774

Table 15	Table 15: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: Criteria Air Pollutants and Female Self-Reported MS cases									
Variable	OR for lowest quartile	OR for second quartile	95% CI <sup>a</sup> of the second quartile OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile's OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the fourth quartile's OR	N (Number of people used for the OR)
PM-2.5 (area)	1.00	0.90	(0.80, 1.03)	1453576	1.02	(0.90, 1.16)	1233432	1.14*	(1.02, 1.28)	2550552
PM-2.5 (all)	1.00	0.84*	(0.71, 1.00)	695928	0.95	(0.82, 1.10)	1229098	1.18*	(1.03, 1.34)	3114398
PM-10 (mobile)	1.00	0.97	(0.78, 1.20)	527312	1.07	(0.89, 1.30)	878612	1.46****	(1.23, 1.74)	3469741
PM-10 (area)	1.00	0.90	(0.79, 1.02)	1256057	0.95	(0.84, 1.08)	1257146	1.12*	(1.00, 1.25)	2758341
PM-10 (all)	1.00	0.89	(0.74, 1.07)	588260	0.88	(0.76, 1.04)	1172605	1.23**	(1.06, 1.42)	3211310
Sulfur dioxide (mobile)	1.00	1.06	(0.87, 1.33)	542359	0.95	(0.77, 1.16)	800885	1.48***	(1.24, 1.77)	3521391
Sulfur dioxide (area)	1.00	0.95	(0.79, 1.16)	633212	1.00	(0.84, 1.19)	870761	1.47***	(1.26, 1.70)	3518047
Sulfur dioxide (all)	1.00	0.93	(0.76, 1.15)	597066	1.10	(0.92, 1.32)	1027098	1.31**	(1.11, 1.56)	3238501

Table 15	Table 15: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: Criteria Air Pollutants and Female Self- Reported MS cases									
				Re	ported MS	cases				
Variable	OR for lowest	OR for second	95% Cl <sup>a</sup> of	N (Number	OR for third	95% CI <sup>a</sup> of the	N (Number	OR for highest	95% CI <sup>a</sup> of the	N (Number of people
	quartile	quartile	the second quartile OR	of people used for the OR)	quartile	third quartile's OR	of people used for the OR)	quartile	fourth quartile's OR	used for the OR)
VOCs (mobile)	1.00	1.00	(0.81, 1.24)	546468	1.10	(0.90, 1.34)	797603	1.41****	(1.19, 1.69)	3514205
VOCs (area)	1.00	0.84	(0.69, 1.03)	549203	0.97	(0.81, 1.16)	868446	1.28**	(1.09, 1.49)	3489057
VOCs (all)	1.00	1.12	(0.90, 1.40)	545179	1.10	(0.89, 1.35)	779424	1.56****	(1.29, 1.88)	3524818

Each variable under analysis included 158 of the 159 counties in GA. VOCs are volatile organic compounds, and PM is particulate matter. Also, the selected air pollutants were reported as emissions in tons, in 1999.

<sup>&</sup>lt;sup>a</sup> 95% CI is calculated using the case-control study estimation model

<sup>\*</sup>Values are significant at the 0.05 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*</sup>Values are significant at the 0.01 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*\*</sup>Values are significant at the 0.001 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*\*\*</sup>Values are significant at the 0.0001 level using a Chi-squared test for 2 x 2 contingency tables

Table 16 shows increasing odds ratios for male MS self-reported cases with increasing emissions of selected criteria air pollutants to outdoor air for PM-2.5 (all sources), sulfur dioxide (all sources), and volatile organic compounds (all sources). In addition, the odds ratios tended to increase from the second to the fourth quartile for nitrogen oxides (mobile and areas sources) and volatile organic compounds (area sources).

Table	Table 16: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: Criteria Air Pollutants and Male Self-Reported MS cases										
Variable	OR for lowest quartile	OR for second quartile	95% CI <sup>a</sup> of the second quartile's OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile's OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the fourth quartile's OR	N (Number of people used for the OR)	
Carbon Monoxide (mobile)	1.00	1.51	(0.98, 2.35)	531636	1.45	(0.96, 2.20)	770204	1.93***	(1.32, 2.85)	3430620	
Carbon monoxide (area)	1.00	1.37*	(1.07, 1.75)	1428572	1.21	(0.93, 1.56)	1173506	1.51***	(1.21, 1.91)	2562559	
Carbon monoxide (all)	1.00	0.76	(0.56, 1.03)	666994	0.73*	(0.56, 0.96)	950091	1.07	(0.86, 1.33)	3381622	
Nitrogen oxides (mobile)	1.00	0.94	(0.65, 1.36)	600947	1.10	(0.79, 1.55)	793340	1.47**	(1.09, 1.99)	3422295	
Nitrogen oxides (area)	1.00	0.99	(0.76, 1.29)	917547	1.09	(0.85, 1.39)	1166838	1.24*	(1.00, 1.54)	3050093	
Nitrogen oxides (all)	1.00	1.15	(0.77, 1.74)	563035	1.14	(0.78, 1.67)	843607	1.74***	(1.23, 2.46)	3354271	
PM-2.5 (mobile)	1.00	1.28	(0.82, 2.02)	495265	1.24	(0.81, 1.90)	750279	1.84**	(1.25, 2.73)	3455027	

Table	16: Odds I	Ratios(OR	) and 95% (		Interval (option	. •	tile: Criteria	Air Pollut	tants and M	ale Self-
Variable	OR for lowest quartile	OR for second quartile	95% CI <sup>a</sup> of the second quartile's OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile's OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the fourth quartile's OR	N (Number of people used for the OR)
PM-2.5 (area)	1.00	1.04	(0.83, 1.32)	1404666	0.96	(0.75, 1.22)	1210591	1.24*	(1.00, 1.54)	2514966
PM-2.5 (all )	1.00	1.16	(0.84, 1.61)	688486	1.22	(0.91, 1.63)	1195959	1.51**	(1.16, 1.98)	3049516
PM-10 (mobile)	1.00	0.96	(0.65, 1.41)	521467	0.95	(0.719, 1.248)	861601	1.415*	(1.121, 1.785)	3388376
PM-10 (area)	1.00	1.04	(0.83, 1.32)	1219051	0.92	(0.72, 1.16)	1235895	1.18	(0.96, 1.45)	2710613
PM-10 (all)	1.00	0.88	(0.63, 1.22)	584408	0.86	(0.65, 1.13)	1139401	1.18	(0.92, 1.53)	3148283
Sulfur Dioxide (mobile)	1.00	1.23	(0.83, 1.85)	534750	1.13	(0.77, 1.65)	797166	1.72**	(1.22, 2.42)	3438621
Sulfur Dioxide (area)	1.00	0.81	(0.58, 1.13)	622797	0.74	(0.54, 1.00)	857461	1.29*	(1.00, 1.66)	3424695
Sulfur Dioxide (all)	1.00	1.05	(0.71, 1.55)	594673	1.32	(0.93, 1.88)	1007391	1.56**	(1.12, 2.18)	3158014

Table	Table 16: Odds Ratios(OR) and 95% Confidence Interval (CI) by Quartile: Criteria Air Pollutants and Male Self- Reported MS cases									
				Rep	orted MS	cases				
Variable	OR for lowest quartile	OR for second quartile	95% CI <sup>a</sup> of the second quartile's OR	N (Number of people used for the OR)	OR for third quartile	95% CI <sup>a</sup> of the third quartile's OR	N (Number of people used for the OR)	OR for highest quartile	95% CI <sup>a</sup> of the fourth quartile's OR	N (Number of people used for the OR)
VOCs (mobile)	1.00	1.51*	(1.00, 2.27)	541492	1.35	(0.92, 1.99)	796058	1.79***	(1.26, 2.56)	3425559
VOCs (area)	1.00	0.68*	(0.48, 0.98)	543221	0.82	(0.60, 1.11)	859140	1.14	(0.87, 1.49)	3399836
VOCs (all)	1.00	1.03	(0.69, 1.53)	542503	1.10	(0.76, 1.59)	767560	1.51*	(1.09, 2.11)	3437659

Each variable under analysis included 158 of the 159 counties in GA. VOCs are volatile organic compounds, and PM is particulate matter. Also, the selected air pollutants were reported as emissions in tons, in 1999.

<sup>&</sup>lt;sup>a</sup> 95% CI is calculated using the case-control study estimation model

<sup>\*</sup>Values are significant at the 0.05 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*</sup>Values are significant at the 0.01 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*\*</sup>Values are significant at the 0.001 level using a Chi-squared test for 2 x 2 contingency tables

<sup>\*\*\*\*</sup>Values are significant at the 0.0001 level using a Chi-squared test for 2 x 2 contingency tables

# V. Discussion and Conclusions

Summary of Figures and Tables

Again, the purpose of this research was to determine the distribution of MS in GA and also to examine the potential relationships between environmental air pollutants and the distribution of MS in GA. From the geographic distribution of MS, we observed the bulk of the total and female self-reported MS prevalence rates seem to center around Atlanta, the largest city in GA. Also, we found the best predictive model for females with MS includes both per-capita income, a characteristic seen in studies done by Mayer and others (Mayer 1981), and PM-10. Males only had per-capita income in their predictive model. In addition, per-capita income seemed to have a positive influence on the total and female rates in the model, while the PM-10 seemed to have a negative influence on these rates in the model. It is important to note this multivariate model only included U.S. census and NEI source emissions data, not TRI source emissions data, because NEI had data for 158 of 159 GA counties. The TRI included data for between 32 and 107 counties, because emissions (total, to specific environmental media, and in processes or waste transported off site) must only be reported by sources to U.S. EPA over certain threshold levels. If nothing in a county emitted enough to be required to report a specific pollutant to TRI, then that county did not have any information on that specific pollutant.

The odds ratios (ORs) of all three population strata (total, female, and male) increased consistently with increases in the total population, male population, and per capita income. Non-consistent increases were observed for source emissions of developmental toxicants, suspected endocrine toxicants, suspected immunotoxicants, and suspected kidney toxicants. The three population strata had lower ORs for people living in the same house in subsequent census collection years.

For source emissions of criteria air pollutants, the ORs for total and female rates increased with increases of carbon monoxide (mobile sources), nitrogen oxides (areas sources), PM-2.5 (mobile sources), PM-10 (mobile sources), and volatile organic compounds (mobile sources). In addition, the ORs increased as we go from the second to the fourth quartile for nitrogen (all sources), PM-2.5 (area and all sources), PM-10 (area and all sources), sulfur dioxide (area and all sources), and volatile organic compounds (area sources). Also, male ORs increased with increasing emissions of PM-2.5 (all sources), sulfur dioxide (all sources), and volatile organic compounds (all sources). Furthermore, the ORs increased from the second to the fourth quartile for nitrogen oxides (mobile and areas sources) and volatile organic compounds (area sources).

Lastly, table 8 showed the percent black and percent white show significantly strong but varying ORs. U.S. Census data does not differentiate Hispanics from the percent black or percent white groups, thus this may partially explain the large jumps in ORs. This may support the need for Census data with percent white and percent black that do not include Hispanic populations.

# Discussion

Prevalence rates help to control for differences in population distribution. Thus, because of the greater prevalence of MS around the largest metropolitan statistical area in GA, Atlanta, and the increasing MS ORs when organized by people per square mile, this study's results suggested a possible role of urban environmental factors in the MS distribution in GA. This coincides with previous studies that suggested that MS rates were positively correlated with per capita income, physicians per capita, steel consumption, and several other characteristics which may increase in cities (Mayer 1981) as well as increased air pollution and other environmental

factors (Oikonen et al, 2003). While this relationship was somewhat weak for the self-reported male prevalence rates, it was stronger for the female prevalence rates.

These differences in the distribution and the models used could be due to several reasons. The majority of MS cases in GA were female. This study has a female to male self-reported case ratio of 3.20 similar to studies conducted by Keegan et al. (2002) and Williamson et al. (Williamson 2007). Thus, in the NMSS-GA data, females were more likely to self-report being diagnosed with MS, females were more likely to become a member, or a combination of both of these characteristics. Thus, the prevalence in females will include a greater number of cases, and thus there may be less internal error in the female prevalence rates than in the male prevalence rates. This could lead to more significant statistics for the females than for the males. Also, there were biochemical gender difference in the mouse model for MS (Spach and Hayes 2005), thus there may also be biochemical gender differences between human MS patients. In addition, while per-capita income has been seen in some studies as associated with MS (Mayer 1981, Catanzaro et al 1992), PM-10 has never before been shown to be associated with MS prevalence rates; however, PM-10 has been shown to be associated with MS relapse in one study conducted in Finland (Oikonen et al. 2003).

Our present study found that self-reported MS prevalence from the National Multiple Sclerosis Society's Georgia Chapter (NMSS-GA) is also related strongly to PM-10 and MS may be positively associated with developmental toxicants, suspected endocrine toxicants, suspected immunotoxicants, suspected kidney toxicants, carbon monoxide (mobile sources), nitrogen oxides (areas and all sources), PM-2.5 (mobile, area, and all sources), PM-10 (mobile, are and all sources), and sulfur dioxide (area and all sources), and volatile organic compounds (area sources).

Biochemically, inhaled particulates like PM-10 can affect a number of host defenses, including the development of specific immune responses such as cell-mediated immunity, the principal part of the immune system implicated in MS. Also, immune stimulation by particulates has also led to increased T-cell activity and immunoglobulin E (IgE) formation. (Bom 2007)

There is a clear relationship, based on many studies published over recent decades, between PM-10 and cardiovascular deaths and respiratory morbidity like asthma symptoms with measured increased outdoor air concentrations of PM-10. Thus, PM-10 may affect not only the cardiovascular and respiratory systems, but also the immune system. Furthermore the immune stimulation and IgE formation could lead to inflammation that is found in MS patients.

In addition to the inflammatory role of PM-10 in MS, heavy metals in particulate matter have been believed to have a relationship with MS clusters (Ingalls 1986; Landrigan et al. 1975). Using odds ratios, we found a significant relationship between lead compounds and female or total MS rates, although this relationship was not seen with males. Nevertheless, this study showed a stronger relationship with particulate matter than the relationship with heavy metals.

The implications of this research are far-reaching. Self-reported MS prevalence was highly associated with PM-10, at the county level, and PM-10 and per-capita income are best associated with MS in our model. Still, the research of environmental air pollutants and MS is sparse, thus additional research is needed to confirm this and the previous study (Oikonen et al. 2003) with richer outcome and exposure data.

Advantages of This Study's Methods

This study was conducted using county level data for several reasons. First, county level data allowed a more intense study of how regional emissions may be associated with regional rates of disease. Also, county level data may allow this study to be repeated in states with

similar data. State emissions data may prove difficult to compare to this study because the distribution of emissions are too large to ensure comparability; however, county level emissions may be specific enough to accomplish this comparison. For example, previous research suggested different latitudes (Mayer 1981; Rosati 2001) made it difficult to compare data with other states further north or south. This study's methodology therefore helped ensure we could compare data from one county to another, instead of comparing GA data with states that have higher or lower MS prevalence because of the effect of sunlight. While this study may have been done at an even smaller level, it was kept at the county level to help ensure continued confidentiality of members of the NMSS-GA.

Also, this study was done using the NMSS-GA member list. To date, a nationwide (beyond NMSS) MS registry has not been created; thus, these data may be from the most accurate database available throughout the country. The NMSS-GA data are also a good source of data, because it is from a national database, and thus the information may be comparable to other NMSS chapter databases throughout the country.

# *Limitations of This Study's Methods*

There are several limitations to this study. First of all, the cases are self-reported from the member's list of the NMSS. While cases were asked to write the date of their MS diagnosis, this information could be fabricated. In addition, there is a small fee to register for the NMSS list. Therefore there is a strong possibility there may be under-reporting and non-differential bias due to information error in our results. Along with these limitations, there may also be relatively more underreporting in males, given generally males are less likely to seek medical care. Still, the NMSS-GA estimated that its list included approximately 75% of the total MS cases in GA.

Also, these data are cross-sectional, although they represented a cumulative prevalence of self-reported MS cases in GA. Therefore, only an estimate of true risk (OR) can be use, and casual inferences cannot be made. Georgia may have different laws, demographics, and other differences from other states. Thus, because this study uses data from GA only, the analyses were interpreted as restricted to GA. There is the possibility these values may be comparable to states with a similar latitude as GA, although different exposure patterns and different environmental laws may make inter-state comparison difficult. Also, birth years of the members were not received, thus this study could not correct for the potential confounding of age. Lastly, using the remaining county population as controls for this study may not be the most robust method for obtaining controls. A better method would be to use those who are also registered as family members of the members of the NMSS-GA, who may be at a higher genetic risk for MS. Conclusions and Recommendations for Future Research and Policy

There may be a role of environmental air pollutants in the start and progression of Multiple Sclerosis (MS). In this study, the best model for predicting the female and total prevalence of MS in GA included both the per-capita income and coarse particulate matter (PM-10). PM-10 along with other environmental air pollutants had highly significant associations with the MS rates in GA, along with evidence of a possible dose response. Moreover, a potential biochemical mechanism does exist. While we do not know the true effects of environmental air pollutants through this study or others, we should explore PM-10 and other air pollutants people are widely exposed to.

It is important to note the exact details of the biochemical mechanism that leads to MS is not currently known. Thus, further study is warranted and desperately needed. Future research should be done using the reported dates of diagnosis, to be able to assess yearly incidence as well

as prevalence. Then, relative risk could be calculated as a true measure of risk, and not OR, which is only an estimate. Also, including the month and year of birth in subsequent studies will facilitate eliminating the ever-present confounding of age. Also, collecting ethnicity-data would be beneficial, because MS has been more common among some races than others (Keegan and Noseworthy 2002; Rosati 2001). This will allow future research to also control for racial differences. Lastly, more robust studies, such as prospective studies that are nested in the past, using various data available, may help provide more robust data for further analysis.

From a policy point of view, a MS registry is desperately needed. While we do have self-reported data that may be somewhat accurate, a universal definition and universal data collection would help scientists solidify the epidemiology of MS and conclusions from these studies. This registry must include important demographic information such as date of birth, race, date of diagnosis by a physician, and county and state of residence. Without a registry, MS data may continue to be underreported. Also, MS research has been supported, and should continue to be supported across the world. This disease has been known for a long time, yet, there is not a concrete list of causal factors. More research may help to bring light and consensus among the many factors that may influence the start and progression of MS, as well as how many people are diagnosed with MS and the common and diverse exposures that they may have.

The distribution and determinants of disease is the foundation of epidemiology. County-level, or lower, graphic representations of MS are important. While MS may seem to be equally distributed throughout the state, it was not in this study. Through finding high-risk counties and determining their differences from low-risk counties, we may be able to gather further crucial information about MS.

As it has for hundreds of years, the question of nature versus nurture rages on. This debate centers on whether genetics or our environment leads more directly to our behaviors and to our diseases. It seems that MS may be a point of synthesis, for it embodies strong genetic and environmental factors. May we work quickly to elucidate this relationship in MS, for lives depend upon it.

# REFERENCES

- Akhyani N, Berti R, Brennan MB, Soldan SS, Eaton JM, McFarland HF, et al. 2000. Tissue distribution and variant characterization of human herpesvirus (HHV)-6: increased prevalence of HHV-6A in patients with multiple sclerosis. J Infect Dis 182(5):1321-1325.
- Anderson DW, Ellenberg JH, Leventhal CM, Reingold SC, Rodriguez M, Silberberg DH. 1992. Revised estimate of the prevalence of multiple sclerosis in the United States. Ann Neurol 31(3):333-336.
- Baum HM, Rothschild BB. 1981. The incidence and prevalence of reported multiple sclerosis. Ann Neurol 10(5):420-428.
- Bolviken B, Celius EG, Nilsen R, Strand T. 2003. Radon: a possible risk factor in multiple sclerosis. Neuroepidemiology 22(1):87-94.
- Bom KEDaP. 2007. Particle Toxicology. Boca Raton, FL:CRC Press.
- Cantorna MT, Hayes CE, DeLuca HF. 1996. 1,25-Dihydroxyvitamin D3 reversibly blocks the progression of relapsing encephalomyelitis, a model of multiple sclerosis. Proc Natl Acad Sci U S A 93(15):7861-7864.
- Cantorna MT, Woodward WD, Hayes CE, DeLuca HF. 1998. 1,25-dihydroxyvitamin D3 is a positive regulator for the two anti-encephalitogenic cytokines TGF-beta 1 and IL-4. J Immunol 160(11):5314-5319.
- Catanzaro M, Weinert C. 1992. Economic status of families living with multiple sclerosis. Int J Rehabil Res 15(3):209-218.
- CDC UatWHO. 2001. Epi Info 6.
- Cepok S, Zhou D, Srivastava R, Nessler S, Stei S, Bussow K, et al. 2005. Identification of Epstein-Barr virus proteins as putative targets of the immune response in multiple sclerosis. J Clin Invest 115(5):1352-1360.
- Dal Canto MC, Melvold RW, Kim BS, Miller SD. 1995. Two models of multiple sclerosis: experimental allergic encephalomyelitis (EAE) and Theiler's murine encephalomyelitis virus (TMEV) infection. A pathological and immunological comparison. Microsc Res Tech 32(3):215-229.
- de la Maza M, Garcia J, Bernal J, Fuentes M. 2000. [A review of the epidemiology of multiple sclerosis in Mexico]. Rev Neurol 31(5):494-495.
- de Rezende PA, Arruda WO. 1996. [Genetic aspects in multiple sclerosis: I. Twin studies]. Arq Neuropsiquiatr 54(3):433-438.
- Dyment DA, Ebers GC, Sadovnick AD. 2004. Genetics of multiple sclerosis. Lancet Neurol 3(2):104-110.
- Ebers GC, Kukay K, Bulman DE, Sadovnick AD, Rice G, Anderson C, et al. 1996. A full genome search in multiple sclerosis. Nat Genet 13(4):472-476.
- Fazekas F, Barkhof F, Filippi M, Grossman RI, Li DK, McDonald WI, et al. 1999. The contribution of magnetic resonance imaging to the diagnosis of multiple sclerosis. Neurology 53(3):448-456.
- Freedman DM, Dosemeci M, Alavanja MC. 2000. Mortality from multiple sclerosis and exposure to residential and occupational solar radiation: a case-control study based on death certificates. Occup Environ Med 57(6):418-421.

- Haahr S, Plesner AM, Vestergaard BF, Hollsberg P. 2004. A role of late Epstein-Barr virus infection in multiple sclerosis. Acta Neurol Scand 109(4):270-275.
- Hansen T, Skytthe A, Stenager E, Petersen HC, Kyvik KO, Bronnum-Hansen H. 2005. Risk for multiple sclerosis in dizygotic and monozygotic twins. Mult Scler 11(5):500-503.
- Henry J, et al. . 2007. Investigation of a Cluster of Multiple Sclerosis in Two Elementary School Cohorts. Journal of Environmental Health 69(10):34-38.
- Hernan MA, Olek MJ, Ascherio A. 2001. Cigarette smoking and incidence of multiple sclerosis. Am J Epidemiol 154(1):69-74.
- Hogancamp WE, Rodriguez M, Weinshenker BG. 1997. The epidemiology of multiple sclerosis. Mayo Clin Proc 72(9):871-878.
- Hong J, Zang YC, Tejada-Simon MV, Li S, Rivera VM, Killian J, et al. 2000. Reactivity and regulatory properties of human anti-idiotypic antibodies induced by T cell vaccination. J Immunol 165(12):6858-6864.
- Ingalls TH. 1986. Endemic clustering of multiple sclerosis in time and place, 1934-1984. Confirmation of a hypothesis. Am J Forensic Med Pathol 7(1):3-8.
- Ito H, Yamasaki K, Kawano Y, Horiuchi I, Yun C, Nishimura Y, et al. 1998. HLA-DP-associated susceptibility to the optico-spinal form of multiple sclerosis in the Japanese. Tissue Antigens 52(2):179-182.
- Keegan BM, Noseworthy JH. 2002. Multiple sclerosis. Annu Rev Med 53:285-302.
- Kong H, Baerbig Q, Duncan L, Shepel N, Mayne M. 2003. Human herpesvirus type 6 indirectly enhances oligodendrocyte cell death. J Neurovirol 9(5):539-550.
- Kurtzke JF. 2000. Multiple sclerosis in time and space--geographic clues to cause. J Neurovirol 6 Suppl 2:S134-140.
- Landrigan PJ, Gehlbach SH, Rosenblum BF, Shoults JM, Candelaria RM, Barthel WF, et al. 1975. Epidemic lead absorption near an ore smelter. The role of particulate lead. N Engl J Med 292(3):123-129.
- Levin LI, Munger KL, Rubertone MV, Peck CA, Lennette ET, Spiegelman D, et al. 2003. Multiple sclerosis and Epstein-Barr virus. Jama 289(12):1533-1536.
- ---. 2005. Temporal relationship between elevation of epstein-barr virus antibody titers and initial onset of neurological symptoms in multiple sclerosis. Jama 293(20):2496-2500.
- Lunemann JD, Edwards N, Muraro PA, Hayashi S, Cohen JI, Munz C, et al. 2006. Increased frequency and broadened specificity of latent EBV nuclear antigen-1-specific T cells in multiple sclerosis. Brain 129(Pt 6):1493-1506.
- Mayer J. 1981. Georgraphical Clues about Multiple Sclerosis. Annals of the Association of American Geographers 71(1):28.
- Microsoft-Corporation. 2004. Microsoft Excel 2004 for Mac.
- Munger KL, Zhang SM, O'Reilly E, Hernan MA, Olek MJ, Willett WC, et al. 2004. Vitamin D intake and incidence of multiple sclerosis. Neurology 62(1):60-65.
- NL Sicotte ea. 2002. Treatment of multiple sclerosis with the pregnancy hormaone estriol. Annuals of Nuerology 52(4):421-428.
- Ogawa G, Mochizuki H, Kanzaki M, Kaida K, Motoyoshi K, Kamakura K. 2004. Seasonal variation of multiple sclerosis exacerbations in Japan. Neurol Sci 24(6):417-419.
- Oikonen M, Laaksonen M, Laippala P, Oksaranta O, Lilius EM, Lindgren S, et al. 2003. Ambient air quality and occurrence of multiple sclerosis relapse. Neuroepidemiology 22(1):95-99.

- Oksenberg JR, Baranzini SE, Barcellos LF, Hauser SL. 2001. Multiple sclerosis: genomic rewards. J Neuroimmunol 113(2):171-184.
- Rosati G. 2001. The prevalence of multiple sclerosis in the world: an update. Neurol Sci 22(2):117-139.
- Soldan SS. 2003. Immune modulation in multiple sclerosis patients treated with the pregnancy hormone estradiol. J Immunol 171(11):6267-6274.
- Soldan SS, Berti R, Salem N, Secchiero P, Flamand L, Calabresi PA, et al. 1997. Association of human herpes virus 6 (HHV-6) with multiple sclerosis: increased IgM response to HHV-6 early antigen and detection of serum HHV-6 DNA. Nat Med 3(12):1394-1397.
- Sospedra M, Martin R. 2005. Immunology of multiple sclerosis. Annu Rev Immunol 23:683-747.
- Sospedra M, Zhao Y, zur Hausen H, Muraro PA, Hamashin C, de Villiers EM, et al. 2005. Recognition of conserved amino acid motifs of common viruses and its role in autoimmunity. PLoS Pathog 1(4):e41.
- Spach KM, Hayes CE. 2005. Vitamin D3 confers protection from autoimmune encephalomyelitis only in female mice. J Immunol 175(6):4119-4126.
- SPSS. 2005. SPSS for Windows.
- Sumaya CV, Myers LW, Ellison GW, Ench Y. 1985. Increased prevalence and titer of Epstein-Barr virus antibodies in patients with multiple sclerosis. Ann Neurol 17(4):371-377.
- Tejada-Simon MV, Zang YC, Hong J, Rivera VM, Zhang JZ. 2003. Cross-reactivity with myelin basic protein and human herpesvirus-6 in multiple sclerosis. Ann Neurol 53(2):189-197.
- Thompson AJ, Polman CH, Miller DH, McDonald WI, Brochet B, Filippi MMX, et al. 1997. Primary progressive multiple sclerosis. Brain 120 ( Pt 6):1085-1096.
- Virtanen JO, Zabriskie JB, Siren V, Friedman JE, Lyons MJ, Edgar M, et al. 2005. Colocalization of human herpes virus 6 and tissue plasminogen activator in multiple sclerosis brain tissue. Med Sci Monit 11d(3):BR84-87.
- Wandinger K, Jabs W, Siekhaus A, Bubel S, Trillenberg P, Wagner H, et al. 2000. Association between clinical disease activity and Epstein-Barr virus reactivation in MS. Neurology 55(2):178-184.
- Weinshenker BG. 1998. The natural history of multiple sclerosis: update 1998. Semin Neurol 18(3):301-307.
- Williamson D, et al. 2007. Prevalence of Multiple Sclerosis in 19 Texas counties, 1998-2000. Journal of Environmental Health 69(10):41-45.
- Williamson DM, Henry JP. 2004. Challenges in addressing community concerns regarding clusters of multiple sclerosis and potential environmental exposures. Neuroepidemiology 23(5):211-216.

Appendices A-C

Appendix A. Selected Variables and Correlations with Multiple Sclerosis Prevalence (caption shows "Table 3") (Mayer 1981)

TABLE 3.—SELECTED VARIABLES AND CORRELATIONS WITH MULTIPLE SCLEROSIS PREVALENCE

Variable	Correlation coefficient	
Mean annual hours of sunshine	88	
Steel consumption (tons/capita)	+.85	
Physicians per capita	+.85	
Literacy rate	+.84	
Latitude (population centroid)	+.77	
Gross energy consumption (megawatt		
hours per capita)	+.76	
Mean annual temperature	70	
Infant mortality	69	
GNP per capita	+.65	
Food consumption (caloric per capita)	+.59	


Source: Leibowitz and Alter, op. cit., footnote 4, p. 261.


Many different factors like mean annual house of sunshine, mean annual temperature and infant mortality were found to be protective of Multiple Sclerosis. Contrarily steel consumption, number of physicians per capita, literacy rate, latitude from equator, gross energy consumption, gross national product per capita and food consumption were all associated with higher rates of MS

(Mayer 1981)

# Appendix B: Figures 1, 2 and 3 (Rosati 2001)

Figures 1, 2 and 3 show Geographical Information Systems maps of Multiple Sclerosis prevalence (per 100,000) in the United States, Africa and around the United Kingdom, respectively. (Rosati 2001)





In all three of these Geographic Information System (GIS) models, Rosati et al. show that there are significant geographical influences that may influence the rate of MS in several countries. (Rosati 2001)

Appendix C: Crude and adjusted odds ratios (95% Confidence Intervals) for mortality from Multiple Sclerosis associated with indicators of exposure to sunlight and socioeconomic status, USA, 1984-1985 (Freedman et al. 2000)

Appendix 3 Crude and adjusted ORs (95% CIs) for mortality from Multiple Sclerosis associated with indicators of exposure to sunlight and socioeconomic status, USA, 1984-

95										
	Multiple Sclerosis				Skin cancer					
	Crude		Adjusted*		Crude		Adjusted			
Residential Sunlight										
Low		1		1		1		1		
Moderate	0.58 (0.54 to		0.59 (0.55 to		1.12 (1.04 to		1.15 (1.07 to			
	0.62)		0.63)		1.20)		1.23)			
High	0.55 (0.51 to		0.53 (0.48 to		1.15 (1.07 to		1.24 (1.15 to			
	0.59)		0.57)		1.24)		1.35)			
Occupational										
Indoor		1		1		1		1		
Mixed	0.87 (0.77 to		0.89 (0.78 to		1.30 (1.19 to		1.03 (0.93 to			
	0.97)		1.02)		1.42)		1.14)			
Outdoor	0.40 (0.33 to		0.74 (0.61 to		1.53 (1.40 to		1.21 (1.09 to			
	0.47)		0.89)		1.68)		1.34)			
Socioeconomic status:										
1 Low		1		1		1		1		
2	1.36 (1.19 to		1.00 (0.88 to		0.92 (0.84 to		0.93 (0.85 to			
	1.55)		1.15)		1.0)		1.01)			
3	2.45 (2.19 to		1.90 (1.68 to		0.90 (0.83 to		0.89 (0.82 to			
	2.75)		2.13)		0.97)		0.96)			
4	3.08 (2.73 to		2.26 (1.99 to	-	0.87 (0.79 to		0.90 (0.82 to			
	3.49)		2.57)		0.95)		1.00)			
5 High	3.39 (2.89 to		3.07 (2.59 to	-	1.19 (1.04 to		1.07 (0.94 to			
	3.99)		3.62)		1.36)		1.23)			

<sup>\*</sup> Adjusted ORs were calculated with logistic regression, adjusting for age, sex, race and socioeconomic status, except in the case of socioeconomic status where adjustments were for age, sex and race.

Residence sunlight was defined as residing in a region both at birth and at the time of death. Regions are defined in Appendix 1. Subjects who resided in different regions at birth and at the time of death were excluded.

Occupational analysis was limited to those subjects whose usual occupation was classified as highly or moderately active to exclude those whose indoor jobs reflected restricted job options due to the disabilities of their disease. Homemakers were also excluded.

Freedman et al. show that residential sunlight is protective of MS (OR=0.53). They also show state that under the highest sun exposure white men, white women, black men and black women (adjusted OR=0.57, 0.49, 0.44, 0.59, respectively[data not shown]) have similar rates. (Freedman et al. 2000)